

Designing an intersectional interventions model for brucellosis occurrence reduction in north-west of Iran

Damari B, MD-PhD¹, Ahmadpour M, MSc^{2*}, Hallajzadeh J, PhD³, Hosseini SH, MSc⁴, Safiri S, PhD⁵, Ghaderpoori M, MSc⁶, Hashemi V, MSc⁷

1- Assistant Prof., National Institute for Health Research, Tehran University of Medical Sciences, Tehran, Iran. 2-Faculty Member, Dept. of Public Health, Maragheh University of Medical Sciences, Maragheh, Iran. 3-Assistant Prof., Dept. of Basic Sciences, Maragheh University of Medical Sciences, Maragheh, Iran. 4-Faculty member, Vector-borne Diseases Research Center, North Khorasan University of Medical Sciences, Bojnurd, Iran. 5-PhD in Epidemiology, Road Traffic Injury Research Center, Dept. of Statistics & Epidemiology, School of Public Health, Tabriz University of Medical Sciences, Tabriz, Iran. 6- Student of PhD, Student Research Office, Dept. of Environmental Health Engineering, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran. PhD in Environmental Health, Faculty of Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran. 7- Faculty Member, Dept. of Basic Sciences, Maragheh University of Medical Sciences, Maragheh, Iran.

Abstract

Received: July 2015, Accepted: August 2015

Background: Brucellosis is a bacterial infection mostly transmitted by animals to humans; thus, consideration of sanitary instances is an important preventive method for this disease. Therefore, we designed a study in order to reduce the rate of brucellosis in Baneh County, Kurdistan Province, Iran.

Materials and Methods: The sources of data were the medical documents of patients diagnosed in health centers and private laboratories during 2011–2012. Data were analyzed using SPSS software, and chi-square and t-test ($P < 0.05$). In addition, in the second section of the research, data on all brucellosis patients and experts were used to design an intersectional preventive interventions model for this county.

Results: A total of 135 brucellosis patients were surveyed in this study during 2011–2012, and a total of 80 experts participated in this research. It was found that 43 patients (31.9%) lived in the city and 92 of them (68.1%) lived in a village; and 71 of them (52%) were men and 64 (48%) were women. The most common way of incurring the disease was drinking non-boiled milk [42 (31.11%)]. Moreover, 100% of the patients and 100% of the experts believed that intersectional intervention is the best prevention method for brucellosis.

Conclusions: Considering the results, it can be concluded that the intersectional intervention with unique management is the best way for reducing the incidence and prevalence rate of brucellosis in Baneh County.

Keywords: Epidemiological Study, Brucellosis, Malta Fever, Iran.

Introduction

Brucellosis is a bacterial zoonotic infection mostly transmitted by animals to humans. It is not transmitted from one human to another, but is a life threatening infection which represents a wide range of clinical diseases in humans mainly caused by *brucella melitensis-abortus* (1-3). Brucellosis was discovered by David Bruce, a Scottish physician, in 1887, and due to the observation of its first case in Malta Island, it has also been named Malta

fever (2-4). In many countries, brucellosis included of reportable diseases. Overall, this disease has unfavorable effects on human and animal health and their products. In animals, this disease is called fetus abortion (3, 5). This disease has other names like Billowy fever, Mediterranean fever, and mad fever. Human

* **Corresponding author:** Mohammad Ahmadpour, Dept. of Public Health, Maragheh University of Medical Sciences, Maragheh, Iran.
Email: moahmadpour@yahoo.com

brucellosis, Malta fever, is a serious public health risk in regions with infectious mutton and goats and seen mostly in spring and summer, the seasons of offspring and suckling of animals (5-7). There are various methods for controlling this disease which are mainly based on prevention. Therefore, surveillance is a key means of management of control, and prevention programs for this disease. Thus, consideration of sanitary instances is an important method for its prevention, and the background of this disease shows that different sectors should be responsible for its incidence and prevalence rate reduction. Based on the official data from the Ministry of Health, Treatment, and Medical Education of Iran, the morbidity rate of brucellosis in Iran, and Baneh County (Kurdistan Province, Iran), respectively, were 11635,354 and 28 cases in 2010 and 14128,536 and 65 cases in 2011. Hence, this data shows that the prevalence rate of brucellosis disease has had an ascending trend in the abovementioned places (7-9). Therefore, the present research was designed to conduct an epidemiological survey and to design an intersectional interventions model for incidence and prevalence rate reduction of brucellosis in Baneh County during 2011-2012.

Material and Methods

This cross-sectional and analytical study is based on several data sources such as all documents in health center and private laboratories in Baneh County and its districts. In addition, a valid and unique questionnaire was used for gathering data. The total number of brucellosis patients surveyed during 2011-2012 in this research was 135 cases. Moreover, the total number of experts who participated in this research was 80

individuals. The demographic and laboratory information of all cases diagnosed during 2011-2012 were studied. These patients were diagnosed using pathological tests in health care centers or private medical centers. Finally, all patients were categorized using a questionnaire based on age, gender, residential area, literacy level, job, the kind of dairy products consumed by the patient, and etcetera. Then, SPSS software (version 19, SPSS Inc., Chicago, IL, USA) and descriptive analysis, chi-square, and t-test were used for the statistical analysis of the cases (P < 0.05). In the second section of the research, the demographic, and viewpoints of all brucellosis patients and experts were collected in order to design an intersectional interventions model to prevent the incidence of and reduce the prevalence rate of brucellosis. In this study, face-to-face and cumulative interviews were performed based on a questionnaire with all experts and patients in Baneh County. Then, through the aggregation of all of this data, an intersectional interventions model was designed. All patients and experts were free to participate in or leave the study. All participants remained in the study.

Results

The demographic data provided in table 1 show that 43 and 92 patients lived in the city and villages, respectively. In addition, 52% of them were men and 48% were women.

The results presented in table 2 illustrate that the most common cause of incurring the disease was drinking non-boiled milk. Furthermore, the majority of cases of morbidity occurred in spring and the lowest rate of morbidity was observed in winter. In addition, 34.07% of patients were ranchers.

Table 1: Frequency distribution of cases with brucellosis in Baneh County according to gender and residential area during 2011-2012

Residential area	Gender				Total	
	Women		Men		N	%
	N	%	N	%		
Village	44	68.75	48	67.60	92	68.10
City	20	31.25	23	32.39	43	31.90

Table 2: Occupational, seasonal, and causal distribution of cases with brucellosis in Baneh County during 2011-2012

	Variable	Number (Percentage) of patients
The Kind of consumed dairy	Nonpasteurized ice-cream	3 (2.22)
	Fresh cheese	36 (26.66)
	Non-boiled milk	42 (31.11)
	Fresh cheese and non-boiled milk	28 (21.59)
	Others	26 (19.25)
Season	Spring	60 (44.40)
	Summer	36 (26.60)
	Autumn	24 (17.70)
	Winter	15 (11.10)
Antecedent of contact with the brucellosis during the last year	Yes	97 (72.30)
	No	17 (12.30)
	Not clear	21 (15.40)
Year	2011	65 (48.20)
	2012	70 (51.80)
Job	Employee	2 (1.48)
	Farmer	17 (12.59)
	Self-employed	5 (3.70)
	Householder wife	20 (20.74)
	Rancher	46 (34.07)
	Student	20 (14.81)
	Labor	8 (5.92)
	Others	9 (6.60)
	Total	135 (100)

Table 3 shows that 86.7% of patients selected the option of “intersectional cooperation under unique management” as the best preventive model for brucellosis.

As table 4 shows, all experts were unanimous in selecting intersectional cooperation as the best method for prevention of brucellosis.

Table 3: Frequency distribution of brucellosis patients based on their viewpoints about brucellosis prevention and intersectional cooperation in Baneh County during 2011-2012

Prevention Method	Number (Percentage)	Organs responsible for brucellosis prevention	Number (Percentage)
Self-centered way	4 (3.0)	Veterinary Office	60 (44.6)
Intersectional cooperation under unique management	117 (86.7)	Agricultural Jihad Office	37 (27.3)
Other	0 (0)	Health Network Office	33 (24.1)
No idea	14 (10.4)	None of them	5 (4.0)
Total	135 (100)	Total	135 (100)

The model was derived in the final step from this research (The model of section1& The model of section2) explains that the intersectional intervention with unique

management is the best way for reducing the incidence and prevalence rate of brucellosis in Baneh County.

Table 4: Frequency distribution of experts based on their viewpoints about intersectional cooperation items for Brucellosis prevention in Baneh County

Intersectional cooperation items which are necessary for Brucellosis prevention	Number (Percentage)
Determination of the Executive Committee with the cooperation of all offices	80 (100)
Activation of the Executive Committee under the Governorship office	80 (100)
Determination of the job description for all offices cooperated	80 (100)
Compelling of all cooperating offices to execute the related affairs	80 (100)
Determination of the Educational Committee by all offices for the education of target groups and the public	80 (100)




Figure 1: Monthly distribution of cases with brucellosis in Baneh County during 2011-2012

The model of section 1: The intersectional interventions model for prevention of brucellosis disease in Baneh County

Administrative structure of the model:

Inviting offices which perform actions relevant to the prevention of brucellosis, including the Governorship Office, Health network Office, Veterinary Office, Agricultural Jihad Office, and Municipal Office.

Explaining the subject at the meeting with the governor and obtaining the signature of the relevant departments
obtaining approval for projects

Formation of the Common Committee for prevention of and fight against brucellosis in Baneh County by the related departments

Formation of the Education Committee with the participation of experts from relevant departments

Production and distribution of a CD and other publications with the subject of brucellosis prevention

Taking advantage of the potential of public participation, NGOs, and city and village councils to prevent the incidence

The Resources needed:

Human resources
Financial resources
Equipment resources
Information resources

The final outcome:

Reduction of the incidence and prevalence rate of brucellosis in Baneh County

The standard of the final outcome:

At least 20% reduction in incidence and prevalence rate of brucellosis compared to 2012 in Baneh County

Costs:

The cost of each of the study stages will be provided by Kurdistan University of Medical Sciences and the Joint Chiefs of prevention and fight against brucellosis in the

and reduce the prevalence rate of brucellosis disease in Baneh County	city of Baneh.
8- Preparation and codification of the program for the prevention and treatment of brucellosis and follow-up by the Common Committee	Execution time :
9- Implementation of the specified tasks by relevant departments	From August 2013 for one year and renewed if necessary
10- Formation of the Monitoring and Evaluation Committee by the relevant departments	End users and stakeholders:
11- Delivery of the monthly report on the progress of the project by the Monitoring and Evaluation Committee	Patients, their families, Veterinary Office, Agriculture Jihad Office, Health Network Office, and the public.

The model of section 2: The intersectional interventions model for prevention of brucellosis disease in Baneh County

Duties related to the model were approved for the related offices.

Description of tasks for the prevention of brucellosis for the related offices in Baneh County	The Name of the related offices
1-Monitoring the implementation of the tasks set for the relevant departments 2- Coordinating relevant departments to carry out their respective duties 3-Monitoring quarterly report measures	Governorship Office
1-Supervision of the production and distribution places of dairy products and ice cream each year, 20% increase in rates compared to last year, and delivery of the quarterly report of actions to the Committee for the prevention of brucellosis 2-Education and empowerment the people to prevent brucellosis disease and provide a 20% reduction in its prevalence compared to the previous year and delivery of the quarterly report of the actions to the Committee for the prevention of brucellosis 3-Case detection and collection of statistical reports on patients with brucellosis, and delivery of the quarterly report of actions to the Committee for the prevention of brucellosis	Health Network Office
1-Planning the vaccination of animals, 20% higher than last year, and delivery of the quarterly report of actions to the Committee for the prevention of brucellosis 2- Prevention of the activity of butchers without a permitted, 20% higher than last year, and delivery the quarterly report of the actions to the Committee for the prevention of brucellosis. 3-The revocation of entry and exit permits for animals in the city, 20% higher than last year, and delivery of the quarterly report of actions to the Committee for the prevention of brucellosis	Veterinary Office
1-Monitoring the maintenance of facilities and animal husbandry, 20% higher than last year, and delivery of the quarterly report of actions to the Committee for the prevention of brucellosis 2-Training ranchers on health procedures of animal husbandry, 20% more than last year, and delivery of the quarterly report of actions to the Committee for the prevention of brucellosis 3-Training ranchers in the field of animal health, milk cattle, and milk transportation and distribution to health practices, 20% more than last year, and delivery of the quarterly report of actions to the Committee for the prevention of brucellosis	Agriculture Jihad Office
Monitoring of animal slaughterhouses, 20% higher than last year, and delivery of the quarterly report to the Committee for the prevention of brucellosis 2-Monitoring of the buying and selling of cattle by local markets in the city, 20% higher than last year, and delivery of the quarterly report to the Committee for the prevention of brucellosis	Municipal Office
Intensifying security measures to prevent illegal trafficking of animals, entering or leaving the boundaries of the city to Iraq, 20% higher than in the past year, and delivery of the quarterly report to the Committee for the prevention of brucellosis	Borderline Guard Office

Discussion

The results of this study show that most cases of brucellosis were observed in 2012 and in

the 2 previous years (2010-2012). Brucellosis had a fluctuating trend. A study performed in East Azerbaijan, Iran, during 8 years (3) and a

study in Arak, Iran, during 10 years showed that the morbidity rate had fluctuated (4). This fluctuation in the morbidity rates can have several causes; for example, inaccuracy in reportage systems, some physicians in certain wards have no commitment to reporting morbidity cases to the responsible offices (5). Moreover, the quality of morbidity reports depends on the related experts following health networks of government wards.

The majority of European countries control and maintain the morbidity trend of brucellosis at a very low level in humans through surveillance systems (6). Based on official data from the Ministry of Health, Treatment, and Medical Education of Iran, the morbidity rate of brucellosis in Iran, and Baneh County were 11635,354 and 28 cases in 2010 and 14128,536 and 65 cases in 2011, respectively. This shows an ascending trend in the prevalence rate of brucellosis (7).

In this study, data were gathered from several data sources based on a valid and unique questionnaire. A total number of 135 brucellosis patients were surveyed during 2011-2012 in this research and a total number of 80 experts participated in this research. The demographic and laboratory information of all cases diagnosed during 2011-2012 were studied. 31.90% and 68.10% of patients lived in the city and villages, respectively. This result is similar to that of the epidemiological study of brucellosis in Maneh and Semelghan, North Khorasan Province, Iran, in 2008-2009 by Shoraka et al.,(8) and the epidemiological study of brucellosis in Kurdistan Province in 2006 by Esmailnasab et al (9). The similarity of morbidity rates in men and women in villages can be due to the similarity in jobs performed by men and women and this is also true about city residents (9, 10).

The incidence of brucellosis in terms of gender was different. the majority of brucellosis cases (52%) were men, and the rest (48%) were women. These results were similar to that of studies conducted in Oromieh, Ardabil, and Kurdistan Provinces, Iran (10). This finding

shows that villagers require more educational programs.

The most common cause of the disease was drinking non-boiled milk (31.11%). Most of the morbidity cases were observed in spring (44.40%) and the lowest rate in winter (11.10%) (11). Furthermore, most brucellosis patients were ranchers (34.07%) (12). These results were similar to that obtained by Esmailnasab et al. and Sofizadeh et al. (13). Nevertheless, in the epidemiological studies of brucellosis by Farahani et al. in Arak, Iran, in 2001-2010 and Soleimani et al. in East Azerbaijan, Iran, in 2001-2009, the most common cause of morbidity was consuming fresh cheese (14). Moreover, these results were similar to that obtained by Brak et al. in their epidemiological study of brucellosis in Ardabil Province, Iran, in 2009 (15). This shows that in order to prevent the incidence of and reduce the rate of brucellosis more educational programs are required.

In this study, most of the patients were ranchers, which is similar to the result of a study performed in Uzbekistan (95.1%) (16). The second highest rate of patients were homemakers (20.74%), which is similar to the result of a study performed in Kurdistan Province (20.8%) (8).

In this study, 44.4% of the morbidity cases occurred in spring, but in the studies conducted in Kashan city and Yazd Province, Iran, most morbidity cases had occurred in the summer (65.5%) (17).

In the present study, 72.3% of the patients had a history of contact with an animal with brucellosis. This rate is higher than that obtained in Yazd Province (63%), and Babol city, Iran, (54.4%) (17).

In our study, 31.11% of the patients had consumed non-boiled (raw) milk and 21.59%, both fresh cheese and non-boiled milk. However, in Babol, 58.1% of patients had consumed milk and cheese (17) and in South Khorasan Province, Iran, 30.2% of the patients had consumed milk or cheese (18).

86.7% of the patients selected the option of “intersectional cooperation under unique

management” as the best preventive method for brucellosis in Baneh County. An example of intersectional cooperation under unique management is all offices, the actions of which are related to the prevention of brucellosis, performing their duty within a common framework provided by the common committee.

We found 100% unanimity among experts regarding intersectional cooperation items for brucellosis reduction in Baneh County.

It can be deduced that intersectional interventions under unique management through consideration of the responsibility of all sectors is the most important way for incidence and prevalence rate reduction of brucellosis. This conclusion is similar to that reached by Mostafavi, Forghani (17), Hatami (18), and Tabatabaei in their studies (19).

Conclusion

It can be concluded that intersectional interventions under unique management through consideration of all the sectors is the most important way for incidence and prevalence rate reduction of brucellosis.

Acknowledgment

We gratefully acknowledge all experts, physicians, and patients who cooperated in this study. We sincerely thank the health personnel of Baneh Health Center for their valuable cooperation in the study.

Conflict of interest: None declared

References

1. Zeynali M, Shirzadi MR, Hajrasoliha H. National Guideline for Brucellosis Control. 2nd ed. Goya MM, Nabavi M, eds. Tehran: Raznahan; 2011.
2. Moradi Gh, Kanani Sh, Majidpour MS, Ghaderi A. Epidemiological status survey of 3880 case of brucellosis in Kurdistan. Iranian Journal of Infectious diseases and Tropical Medicine 2006; 11(33):27-33.
3. Soleimani A , Alizadeh S , Seaif Farshad M, Kusha A, Mohamzadeh M, Haghiri L & et.al. Descriptive Epidemiology of Human Brucellosis

- in East Azerbaijan, 2001-2009. Medical Journal of Tabriz University of Medical Sciences & Health Services 2012; 34(1):63-9.
4. Almasi-Hashiani A, Khodayari M, Eshtrati B, Shamsi M. Factors affecting the interval between the onset and diagnosis of brucellosis in Markazi Province, Iran (2010-11). Arak University of Medical Sciences Journal 2012; 14(7):21-30.
5. Shoraka HR, Hosseini SH, Sofizadeh A, Avaznia A, Rajabzadeh R, Hejazi A. Epidemiological Study of brucellosis in Maneh & Semelghan town, North Khorasan Province in 2008-2009. Journal of North Khorasan University of Medical Sciences 2010; 2(2-3):65-72.
6. Motlagh ME, Oleyaemanesh AR, Beheshtean M. Health and social determinants of it. 2nd. Tehran: Movafagh; 2008.
7. Maleki F, RafieManesh H, Abbasi-Ghahramanloo A, Ghadimi N, Kousha A, Safiri S. Epidemiological characteristics of human brucellosis in Shahin Dezh, Western Azarbaijan, Iran, 2008-2012. Archives of Clinical Infectious Diseases 2015; 10(1):1-4.
8. Esmaeilnasab N, Banafshi O, Ghaderi E, Bidarpour F. Epidemiological change investigation of brucellosis in Kurdistan province in 2006-2007. J Vet Med 2007; 1(3):53-8.
9. Sofizadeh A, Ghorbani M, Salehi R, Mansorian M. Epidemiological study of brucellosis in kalaleh district, Golestan province, in 2003-2007. Journal of Research Development in Nursing & Midwifery 2008; 5(2):8-1.
10. Mantur BG, Amarnath SK, Shinde RS. Review of clinical and laboratory features of human brucellosis. Indian J Med Microbiol 2007; 25(3):188-202.
11. Farahani Sh, Shahmohamadi S, Navidi I, Sofian M. An investigation of the epidemiology of brucellosis in Arak City, Iran 2001-2010. Arak Medical University Journal 2010; 14(6, Suppl 3):49-54.
12. Aghakhani N, saatsaz S, Sharifinia H, Soltani B, Rahbar N, Tabari F. Brucellosis epidemiology in Oromieh city. Poyesh 2012; 3:9-10.
13. Brak M, Emdadi D, Zakipor GH, Seyfnejad SH, Modaressadrdani N. Epidemiological Study of brucellosis in Ardabil Province in 2009. Paper presented at: The National Congress of Brucellosis; 2011 Sep 8-9; Arak, Iran.
14. Mahbobi S, Dorodgar A, Nematian M. Six Years Brucellosis Aspect in the district of Kashan, 2000-2005. Paper presented at: The 2nd National Iranian Congress on Brucellosis; 2007 May 19-21; Tehran, Iran.
15. Karami M, Akbarzadeh Pasha H, Moudi S, ShaHabandaz H, Ghanbari M, Khalilpur A, et al. An epidemiologic study of brucellosis in Babylon city in 6 years. Paper presented at: The 2nd

- National Iranian Congress on Brucellosis; 2007 May 19-21; Tehran, Iran.
16. Mostafavi E, Asmand M. Trends of Brucellosis (Malta fever) in Iran during the period 1991-2008. *Iranian Journal of Epidemiology* 2012; 8(1):94-101.
 17. Earhart K, Vafakolov S, Yarmohamedova N, Michael A, Tjaden J, Soliman A. Risk factors for brucellosis in Samarqand Oblast, Uzbekistan. *Int J Infect Dis* 2009; 13(6):749-53.
 18. Forghani H, Nezamolhoseani SM. Investigate its epidemiology of brucellosis in Yazd province, 2003- 2006. Paper presented at: The 2nd National Iranian Congress on Brucellosis; 2007 May 19-21; Tehran, Iran.
 19. Hatami H. Brucellosis epidemiology. Paper presented at: The 2nd National Iranian Congress on Brucellosis; 2007 May 19-21; Tehran, Iran.
 20. Tabatabaei SM, Zahraei M, Ahmadnai H, Ghotbi M, Rahimi F. Principles of prevention and care of diseases. 2nd. Guya MM, ed. Tehran: Roohghalam; 2007.