



Prevalence and Determinants of Hypertension among Iranian Taxi Drivers (2018)

Payam Khanlari¹, Masoud Khosravipour^{2*}, Faramarz Gharagozlou³, Rashid Heidarimoghadam⁴, Mohammad Babamiri⁵

1- MSc Student, Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran.

2- MSc Student, Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran.

3- Assistant Prof., Dept. of Occupational Health Engineering, School of Health, Kermanshah University of Medical Sciences, Kermanshah, Iran.

4- Professor, Dept. of Ergonomics, School of Health, Hamadan University of Medical Sciences, Hamadan, Iran.

5- Assistant Prof., Dept. of Ergonomics, School of Health, Hamadan University of Medical Sciences, Hamadan, Iran.



Citation: Khanlari P, Khosravipour M, Gharagozlou F, Heidarimoghadam R, Babamiri M. Prevalence and Determinants of Hypertension among Iranian Taxi Drivers in 2018. *JOHE* 2020; 9(1):1-9.

Article Info

* Corresponding author:

Masoud Khosravipour,


E-mail:

masoudkhosravipour74@gmail.com

Article history

Received: Jan 2020

Accepted: Aug 2020

 10.29252/johe.9.1.1

Print ISSN: 2251-8096

Online ISSN: 2252-0902

Peer review under responsibility of Journal of Occupational Health and Epidemiology

Abstract

Background: The prevalence of hypertension (HTN) among Iranian taxi drivers has not been well defined. This study aimed to determine the prevalence of HTN and to identify factors influencing its prevalence among Iranian taxi drivers in 2018.

Materials and Methods: This cross-sectional study was conducted on a total sample of 321 taxi drivers in Kermanshah province in 2018. The participants were classified into two groups, including the HTN group (SBP \geq 140 or DBP \geq 90) and the non-HTN group (SBP $<$ 140 or DBP $<$ 90). A self-administrated questionnaire, including questions about demographic information, medical history, and work-related factors was employed to collect data.

Results: According to the observations, 59.8% of the participants (192/321) suffered from HTN. According to the adjusted logistic regression model and the per one unit increase, there was significantly higher odds of HTN for age (odds ratio [OR] =1.04 and 95% confidence interval [CI] =1.02, 1.06) and work history (OR=1.08 and 95% CI=1.03, 1.13); in contrast, there was a lower risk of HTN for sleep duration (OR=0.83 and 95% CI=0.69, 0.99) and Work Days Per Week (WDPW) (OR=0.73 and 95% CI=0.57, 0.94). In the categorical analysis, a significantly higher risk of HTN was observed in the drivers with age \geq 45, work history \geq 5, Work Hours Per Day (WHPD) $>$ 10, WDPW \leq 6, married, and poor economic status \leq a 2-million Toman income per month.

Conclusions: This study introduces taxi drivers as a high-risk HTN group in the society. Hence, they should be informed of HTN risk factors and undergo regular checkups.

Keywords: Hypertension, Cardiovascular Diseases, Risk Factors, Iran.

Introduction

Hypertension (HTN) is well established as the most common chronic disease worldwide. It is predicted that the total number of hypertensive patients will rise from 972 million individuals in 2000 to 1.56 million individuals in 2025 [1]. Since HTN is asymptomatic in acute stages, the morbidity and mortality resulting from it could be

dramatically high [2]. Approximately 47% of ischemic heart diseases and 57% of stroke cases, being among the prime causes of death in the world, are related to HTN [3-5]. Besides, about 9.5 million annual deaths (16.5% of the total deaths) are attributed to HTN [6]. It was observed that the incidence and mortality of kidney diseases caused by HTN in 2007 and 2017 significantly increased by 28% and 41.4%, respectively [5, 7]. Hence,

identification of high-risk subjects and risk factors affecting HTN prevalence is a high priority.

Professional driving puts drivers under stressful conditions, such as a sedentary lifestyle, an unhealthy diet plan, irregular work schedules [8-10], external stressors, such as noise [11], vibration [12], air pollution [13, 14], and toxic vehicle exhaust emissions [15]. Exposure to these factors turns professional drivers, including bus, truck, and taxi drivers, into high-risk groups for cardiovascular diseases (CVDs). In the past few decades, some studies investigated CVDs among professional drivers. The results of a cohort study, which investigated hospital admissions among professional drivers in Denmark, indicated that age-standardized hospital admissions ratios (SHRs) for CVDs, including ischemic heart disease, myocardial infarction, and cerebrovascular disease, were significantly higher in male taxi and bus drivers than in other drivers [16]. Another study indicated a stronger risk of myocardial infarction among drivers [17]. Furthermore, in another study, a higher risk of stroke was reported in drivers than in the general population [18]. Similarly, in a series of other studies which investigated cardiovascular risk factors, greater risks of metabolic syndrome [19-22], poor sleep quantity and quality [23-26], as well as HTN [27-36] were reported among professional drivers.

Up until now, although many studies have investigated the prevalence of HTN among professional drivers, including bus and truck drivers, little evidence exists on investigating the prevalence of HTN among taxi drivers. In addition, few studies have examined the risk factors influencing the prevalence of HTN among drivers, and in particular, among taxi drivers. To the best of our knowledge, no study has investigated this association among Iranian taxi drivers. Therefore, this study aims to fill this gap by determining the prevalence of HTN and identifying its determinants among Iranian taxi drivers.

Materials and Methods

This cross-sectional study was carried out in Kermanshah province located in western Iran in 2018 in the three months of spring to early summer. Multistage sampling was used to select the study participants. To this end, Kermanshah province was divided into the five central, western, eastern, southern, and northern areas (strata). Next, a similar number of participants were selected from each study area (about 74 drivers in each area). We determined the sample size according to formula 1 as follows:

$$n = \frac{z^2 p (1-p)}{d^2}$$

[1]

Where n indicates the required sample size at the 95% level of significance, p represents the prevalence of HTN among the taxi drivers, which is set at 0.4 according to the previous studies [8, 27, 29-30, 38-41], and d is the degree of precision set at 0.05. Accordingly, the sample size consisted of 369 subjects. In the present study, we only included male taxi drivers licensed by the Taxi Organization, so private taxi drivers were not included. After data collection, taxi drivers with less than a one-year work experience were excluded. In the end, the data obtained from 321 taxi drivers were analyzed.

For the purpose of data collection, a two part self-administrated questionnaire, including questions about demographic information and work-related factors, was employed. Demographic data included age, height, weight, marital status (married or single), current smoking habits, education levels (high school diplomas, lower degrees, and university degrees), economic status (poor status was \leq a 2-million Toman income per month, medium status was between a 2- and a 3-million Toman income per month, and good status was $>$ a 3-million Toman income per month), as well as self-reported diseases and familial medical history. Besides, work-related factors included work experience, work hours per day (WHPD), work days per week (WDPW), self-reported fatigue (SRF), and sleep duration. We used an analog scale for measuring SRF, which ranged from 0 (no fatigue) to 100 (the highest level of fatigue). Sleep duration was determined for each subject by questions "at what hours at night did you usually go to sleep and at what hours did you wake up in the past months?" Moreover, a body mass index (BMI) was calculated for each participant via dividing weight (kg) by height (m²). In addition, the participants' blood pressure was measured by a Beurer BC16 Blood Pressure Monitor according to the standard protocol [37]. In this study, HTN was defined as systolic blood pressure (SBP) \geq 140 mmHg or diastolic blood pressure (DBP) \geq 90 mmHg, according to the latest definition provided by the European Society of Cardiology/the European Society of Hypertension (ESC/ESH) [37]. It is worth noting, to minimize the measurement errors, blood pressure was measured again at a 10-minute interval for each participant, with the average blood pressure computed and recorded in the end.

After importing the data to SPSS software version 25 and classifying them, the normality of the distribution of the quantitative variables was determined by the Shapiro-Wilk test. Accordingly,

the Mann–Whitney test was utilized to assess the significance of the difference between the study groups. In addition, to investigate the significant difference between the study groups for nominal variables, the Chi-square test was utilized. In this study, quantitative and nominal variables were represented by means (the standard deviation) and numbers (percentages). In addition, logistic regression was utilized to examine the effects of study variables on the risk of HTN. Besides, logistic regression was presented by the odds ratio (OR) at a 95% confidence interval (CI) for both unadjusted and adjusted models. It is worth noting that the Spearman's correlation coefficient was employed to examine correlations between the study variables before including them in the model. Moreover, in the adjusted logistic regression model, those variables were considered, which

were significantly different among the study groups, or which had a significant association with HTN (p -value < 0.10) in the simple logistic regression.

Results

According to the study results, 59.8% of the participants (192/321) suffered from HTN. The summary of the study groups' characteristics, including demographic data, work-related factors, as well as self and family medical histories have been provided in Table 1. The results indicate a significant difference between the two groups only in the variables of age, work history, WDPW, marital status, education levels, and self-reported economic status ($p < 0.05$).

Table 1. The summary of the characteristics of the included taxi drivers, Kermanshah province, 2018

Variables		Total subjects (n=321)	Study groups		P-value
			NHTN (n=129)	HTN (n=192)	
			Mean (SD)/ N (%)	Mean (SD)/ N (%)	
SBP, mmHg		137.3 (21.52)	120.1 (10.8)	148.9 (19.0)	-
DBP, mmHg		86.0 (11.7)	78.3 (7.1)	91.2 (11.3)	-
Age, (year)		43.8 (12.1)	40.7 (10.9)	45.9 (12.4)	$< 0.001^a$
BMI, (kg/m ²)		25.1 (3.0)	25.2 (2.9)	25.0 (3.1)	0.452 ^a
Work history, (year)		7.3 (8.3)	4.6 (4.8)	9.1 (9.6)	$< 0.001^a$
WHPD		8.6 (2.1)	8.2 (1.7)	8.8 (2.3)	0.051 ^a
WDPW		5.9 (1.0)	6.0 (1.0)	5.8 (1.1)	0.030 ^a
SRF, (%)		69 (20.0)	70.0 (20.6)	68.2 (19.7)	0.583 ^a
Sleep duration, (hour)		7.0 (1.3)	7.2 (1.3)	6.9 (1.2)	0.088 ^a
Current smoking	Yes	128 (39.9)	54 (41.9)	74 (38.5)	0.552 ^b
Marital status	Married	272 (84.7)	99 (76.7)	173 (90.1)	0.001 ^b
Regular exercise	Yes	70 (21.8)	33 (25.6)	37 (19.3)	0.179 ^b
Education	University degree	54 (16.8)	32 (24.8)	22 (11.5)	0.002 ^b
	High school diploma and less	267 (83.2)	97 (75.2)	170 (88.5)	
Economic status, income in a month	Poor, (≤ 2 million Tomans)	148 (46.1)	47 (36.4)	101 (52.6)	0.004 ^b
	Medium and good (> 2 million Tomans)	173 (53.9)	82 (63.6)	91 (47.4)	
Previous diagnosed diseases	HTN, yes	88 (27.4)	29 (22.5)	59 (30.7)	0.104
	Kidney, yes	67 (20.9)	29 (22.5)	38 (19.8)	0.561 ^b
	Diabetes, yes	72 (22.4)	27 (20.9)	45 (23.4)	0.598 ^b
	Heart, yes	57 (17.8)	21 (16.3)	36 (18.8)	0.570 ^b
Family disease history	HTN, yes	131 (40.8)	46 (35.7)	85 (44.3)	0.124 ^b
	Kidney, yes	112 (34.9)	45 (34.9)	67 (34.9)	0.998 ^b
	Diabetes, yes	97 (30.2)	39 (30.2)	58 (30.2)	0.996 ^b
	Heart, yes	102 (31.8)	43 (33.3)	59 (30.7)	0.623 ^b

Abbreviations: HTN: hypertension; NHTN: non-hypertensive; SD: standard deviation; N, number; BMI: body mass index; WHPD: work hours per day; WDPW: work days per week; SRF: self-reported fatigue; SBP: systolic blood pressure; DBP: diastolic blood pressure
^a Mann–Whitney test for the difference between the two groups; ^b Chi-square test for the difference between the two groups;
 A difference is significant at p -value < 0.05 .

The results of the logistic regression are presented in Table 2. In both adjusted and unadjusted models, it was observed that the subjects aged ≥ 45 years, married, and with poor economic status had a significantly higher risk of HTN than their reference group. In addition, in terms of the

variable of educational level, there was a significantly higher risk of HTN in the subjects with a high school diploma and less, than the reference group in the unadjusted model at p-value < 0.05 and in the adjusted model at p-value < 0.10 .

Table 2. The logistic regression models for hypertension in the included taxi drivers according to demographic and medical history data, Kermanshah province, 2018

Variables	No (%) HTN/NHTN	Logistic regression models				
		Simple		Multiple		
		OR (95%CI)	p-value	OR (95%CI)	p-value	
Age ^a	< 45	97(50.5)/82(63.6)	Referent	-	Referent	-
	≥ 45	95(49.5)/47(36.4)	1.71(1.08,2.70)	0.022	1.68(1.04,2.76)	0.035
BMI ^b	< 25	59(41.3)/70(39.3)	Referent	-	Referent	-
	≥ 25	84(58.7)/108(60.7)	0.92(0.59,1.44)	0.726	0.90(0.56,1.46)	0.672
Marital status ^a	Single	19(38.8)/30(61.2)	Referent	-	Referent	-
	Married	173(63.3)/99(36.4)	2.76(1.48,5.16)	0.001	2.57(1.32,5.01)	0.006
Regular exercise ^b	Yes	37(52.9)/33(47.1)	Referent	-	Referent	-
	No	155(61.8)/96(38.2)	1.44(0.84,2.46)	0.181	1.21(0.68,2.15)	0.517
Smoking habit ^b	No	118(61.1)/75(38.9)	Referent	-	Referent	-
	Yes	74(57.8)/54(42.2)	0.87(0.55,1.37)	0.552	0.91(0.56,1.47)	0.685
Education level ^c	University	22(11.5)/32(24.8)	Referent	-	Referent	-
	High school diploma and less	170(88.5)/97(75.2)	2.55(1.40,4.63)	0.002	1.73(0.92,3.28)	0.092
Economic status ^b	Good	91(47.4)/82(63.6)	Referent	-	Referent	-
	Poor	101(52.6)/47(36.4)	1.94(1.23,3.06)	0.005	1.72(1.07,2.78)	0.026
Previous diagnosed disease^b						
HTN	No	133(69.3)/100(77.5)	Referent	-	Referent	-
	Yes	59(30.7)/29(22.5)	1.53(0.91,2.56)	0.105	1.35(0.77,2.37)	0.293
Kidney	No	154(60.6)/100(39.4)	Referent	-	Referent	-
	Yes	38(56.7)/29(43.3)	0.85(0.49,1.47)	0.561	0.78(0.44,1.39)	0.395
Diabetes	No	147(59.0)/102(41.0)	Referent	-	Referent	-
	Yes	45(62.5)/27(37.5)	1.16(0.67,1.98)	0.598	0.90(0.50,1.61)	0.713
Heart	No	156(59.1)/108(40.9)	Referent	-	Referent	-
	Yes	36(63.2)/21(36.8)	1.19(0.66,2.14)	0.570	1.18(0.62,2.24)	0.610
Family medical history^b						
HTN	No	107(56.3)/83(43.7)	Referent	-	Referent	-
	Yes	85(64.9)/46(35.1)	1.43(0.91,2.67)	0.124	1.19(0.72,1.94)	0.499
Kidney	No	125(59.8)/84(40.2)	Referent	-	Referent	-
	Yes	67(59.8)/45(40.2)	1.00(0.63,1.60)	0.998	0.83(0.50,1.38)	0.471
Diabetes	No	134(59.8)/90(40.2)	Referent	-	Referent	-
	Yes	58(59.8)/39(40.2)	1.00(0.61,1.62)	0.996	0.88(0.52,1.48)	0.627
Heart	No	133(60.7)/86(39.3)	Referent	-	Referent	-
	Yes	59(57.8)/43(42.2)	0.89(0.55,1.43)	0.623	0.75(0.45,1.25)	0.264

Abbreviations: HTN: hypertension; NHTN: non-hypertensive; OR: odds ratio; CI: confidence interval; BMI: body mass index

^aWHPD, WDPW, sleep duration, and economic status

^bWHPD, WDPW, sleep duration, economic status, and age

^cWHPD, WDPW, sleep duration, economic status, and work history

An association is significant at p-value < 0.05 .

The effects of work-related factors, including work history, WHPD, WDPW, SRF, and sleep duration on the risk of HTN have been presented in Table

3. We observed that drivers with a 5-year experience and higher who worked 10 hours and more per day had a significantly higher risk of HTN

than the reference group. Interestingly, there was a lower risk of HTN in the subjects who worked 7

days per week than those who worked 6 days and less per week.

Table 3. The logistic regression models for hypertension in the included taxi drivers according to work-related factors, Kermanshah province, 2018

Variables	No (%) HTN vs. NHTN	Logistic regression models				
		Simple		Multiple		
		OR (95% CI)	P-value	OR (95% CI)	P-value	
Work history ^a	< 5	84(43.8)/89(69.0)	Referent	-	Referent	-
	≥ 5	108(56.3)/40(31.0)	2.86(1.79,4.58)	< 0.001	2.20(1.34,3.63)	0.002
WHPD ^b	≤ 8	97(50.5)/75(58.1)	Referent	-	Referent	-
	9-10	65(33.9)/47(36.4)	1.08(0.66,1.73)	0.785	1.08(0.65,1.81)	0.763
	≥ 10	30(15.6)/7(5.4)	3.31(1.38,7.96)	0.007	2.64(1.02,6.81)	0.045
WDPW ^c	≤ 6	134(69.8)/77(59.7)	Referent	-	Referent	-
	7	58(30.2)/52(40.3)	0.64(0.40,1.23)	0.062	0.55(0.33,0.90)	0.018
SRF ^d	< 60	78(40.6)/58(45.0)	Referent	-	Referent	-
	≥ 60	114(59.4)/71(55.0)	1.19(0.76,1.87)	0.441	1.23(0.76,1.99)	0.390
Sleep duration ^d	≥ 7	69(35.9)/39(30.2)	Referent	-	Referent	-
	≤ 6	123(64.1)/90(69.8)	1.30(0.80,2.09)	0.289	1.43(0.86,2.38)	0.167

Abbreviations: HTN: hypertension; NHTN: non-hypertensive; OR: odds ratio; CI: confidence interval; BMI: body mass index; WHPD: work hours per day; WDPW: work days per week; SRF: self-reported fatigue; HTN: hypertension; NBP: normal blood pressure

^aWHPD, WDPW, sleep duration, economic status, and education levels

^bWDPW, sleep duration, economic status, and age

^cWHPD, sleep duration, economic status, and age

^dWHPD, WDPW, sleep duration, economic status, and age

An association is significant at p-value < 0.05.

Table 4 shows the unadjusted and adjusted logistic regression models for the estimation of the risk of HTN in the case of per unit increase in the quantitative variables. In both models, upon a one-year increase in the work history, the risk of HTN increased significantly. However, it was observed that upon a one-day increase in WDPW, the risk of HTN decreased significantly in both models.

According to the unadjusted model, there was a significantly higher risk of HTN in terms of the variables of WHPD and sleep duration. In contrast, in accordance with the adjusted model, the risk was significant at p<0.10. In terms of other variables, including BMI and SRF, no significant association was observed.

Table 4. The logistic regression models for hypertension per 1-fold increase in the quantitative variables among taxi drivers, Kermanshah province, 2018

Variables	Logistic regression models			
	Simple		Multiple	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Age ^a	1.04(1.02,1.06)	< 0.001	1.04(1.02,1.06)	0.001
BMI ^b	0.99(0.91,1.06)	0.696	0.97(0.90,1.05)	0.512
Work history ^c	1.10(1.05,1.14)	< 0.001	1.08(1.03,1.13)	< 0.001
WHPD ^d	1.14(1.02,1.27)	0.023	1.11(0.98,1.26)	0.095
WDPW ^e	0.77(0.62,0.97)	0.024	0.73(0.57,0.94)	0.013
Sleep duration ^b	0.85(0.71,1.1)	0.069	0.83(0.69,0.99)	0.050
SRF ^b	1.00(0.99,1.01)	0.599	1.00(0.99,1.01)	0.779

Abbreviations: OR: odds ratio; CI: confidence interval; BMI: body mass index; WHPD: work hours per day, WDPW: work days per week; SRF: self-reported fatigue.

^aWHPD, WDPW, sleep duration, and economic status

^bWHPD, WDPW, sleep duration, economic status, and age

^cWHPD, WDPW, sleep duration, economic status, and education levels

^dWDPW, sleep duration, economic status, and age

^eWHPD, sleep duration, economic status, and age

An association is significant at p-value < 0.05.

Discussion

One of the objectives of this study was to determine the prevalence of HTN among Iranian taxi drivers, which had not been investigated in the past. It was observed that the prevalence of HTN was significantly high among Iranian taxi drivers. Accordingly, more than half of the participants suffered from HTN (59.8%). In this study, a higher prevalence of HTN was observed than in previous published studies that reported the prevalence of HTN within the range of 18.2% and 57% [8, 27, 29-30, 38-41]. Furthermore, the comparison of the results of the present study with other studies that investigated the prevalence of HTN among other professional drivers, including suburban, bus, and truck drivers, showed that the prevalence of HTN was significantly higher in taxi drivers than in other professional drivers [28, 32-33, 42-46]. There are several reasons for the higher prevalence of HTN among taxi drivers, among which one could refer to simultaneous exposure to several risk factors, such as a sedentary lifestyle, an unhealthy diet plan, irregular work schedules [8-10], as well as external stressors, such as noise [11], vibration [12], air pollution [13, 14], and toxic vehicle exhaust emissions [15].

Another purpose of this study was to determine the factors probably playing a significant role in the prevalence of HTN among taxi drivers. To this end, we collected data from different lifestyle, medical, and work-related factors and used them in the multiple logistic regression models to examine the association between these factors and HTN. According to the results, a one-year increase in the participants' age significantly increased the risk of HTN by 4%. According to the categorical analysis, subjects aged ≥ 45 had a significantly higher risk of 68% in HTN than subjects aged < 45 . These findings were similar to those of most previous studies on professional drivers [30, 33-35, 41, 43]. However, some studies show no significant association between the drivers' age and the prevalence of HTN [36, 38]. Nevertheless, aging is a well-established risk factor for HTN. Aging, through several biological pathways, including inflammation, oxidative stress, and endothelial dysfunction, could lead to an increased risk of HTN [47]. In this study, a significant association was established between work history and HTN. Accordingly, there was a higher risk of HTN by 8%, per one year increase in work history. Besides, the risk of HTN increased significantly by 120% in the subjects with work history ≥ 5 years versus work history < 5 years. This finding was in line with previous studies that reported an increased risk of HTN with a rise in work history [27, 30, 34, 41, 43].

It is worth noting that due to the high correlation between age and work history, the variable of age played an interfering role in the association.

In addition, we investigated several work-related risk factors, including WDPW, WHPD, SRF, and sleep duration. According to the findings from this research, taxi drivers working more than 10 hours per day were 2.64 times more at the risk of HTN than the reference group (workers who worked 8 hours and less per day). In addition, there was a significantly higher risk of HTN upon an hour increase in WHPD (at p -value < 0.10), which was due to long sedentary times and less physical activities among the drivers. It is well established that low levels of physical activity or long sedentary times could significantly raise the risk of HTN. Moreover, taxi drivers working more hours per day could be more exposed to other parameters associated with HTN. According to some studies, exposure to traffic noise and air pollution could significantly raise the risk of HTN as well. In addition, drivers working more than 10 hours per day reported shorter sleep durations per night than drivers working 9-10 hours per day (6.4 vs. 7.2 h) as well as drivers who worked eight hours and less per day (6.4 vs. 7.1 h). Short sleep durations could be associated with a higher risk of HTN. In addition, a meta-analysis reported a significant dose-response relationship between sleep duration and HTN [26].

Interestingly, we noticed a lower risk of HTN (45%) in the drivers working all days per week than the workers who had at least one day off. In contrast, upon a one-day increase in the working days, the risk of HTN decreased by 27%. Although the two groups had no significant differences in age, BMI, SRF, and sleep duration, the drivers working all days per week had averagely 2 years of lower work history than the other group. In this study, the participants' sleep durations were examined as well, according to which, there was no significant association between the study groups in terms of sleep duration. In addition, it was observed that upon a one-hour increase in sleep duration, the risk of HTN significantly declined by 17%. A previous meta-analysis reported a significant dose-response relationship between sleep duration and HTN [26]. However, no significant association was observed between SRF status and prevalence of HTN among taxi drivers in this study.

In terms of lifestyle factors, a significant association was observed among marital status, economic status (p -value < 0.05), and education levels (p -value < 0.10). In addition, married taxi drivers were shown to be 2.57 times more than others at the risk of HTN, which was consistent with some previous studies [30, 33]. Accordingly,

this finding could be due to the higher mean age (46.4 vs. 29.5), longer work history (8.1 vs. 2.7), and higher BMI (25.2 vs. 24.3) in the married drivers than in the single ones. However, some studies did not show a significant association between marital status and HTN [38, 43]. In terms of the variable of economic status, there was a higher risk of HTN by 72% in the drivers who reported poor economic status than in the reference group. This finding was consistent with a study that investigated this relationship among bus drivers in India [43]. However, it was inconsistent with other studies that showed no significant difference between hypertensive and non-hypertensive drivers in terms of economic status [30, 33, 34]. It was also observed, in this study, that drivers with lower educational levels (high school diplomas and less) had a significantly higher risk of HTN (by 73%) than subjects with university diplomas, at the significance level of $p < 0.10$. This finding was consistent with a study conducted on 491 taxi drivers in Brazil [41]. Although some studies did not show a significant association between educational levels and HTN [33, 34, 43], a recent meta-analysis reported that lower educational levels, as an independent risk factor, increased the risk of HTN (pooled OR= 2.02 and 95% CI= 1.55–2.63) [49]. In the present study, no significant association was found between the variables of BMI, smoking habit, physical activity, medical history, and HTN. These findings were similar to the previous studies on BMI [36], smoking habits [30, 33-36, 50], physical activity [33, 38, 50], medical history and HTN [36], diabetes [33, 36], family HTN [33], and family diabetes [33].

This study was faced with several limitations. Therefore, the results should be interpreted cautiously. Firstly, a cross-sectional design has some limitations in showing cause-effect relationships. Secondly, the use of a convenience sampling method with 321 drivers is not representative of the entire taxi driver population. Thirdly, we were not able to investigate the participants' diet plans. Finally, some of the variables in the present study, including sleep duration and SRF, were collected by subjective methods that could have been associated with measurement errors.

Conclusion

This study was the first attempt to determine the prevalence of HTN among Iranian taxi drivers. In addition, it was a unique attempt to identify risk factors playing a significant role in the prevalence of HTN in the taxi drivers. The findings obtained

consider taxi drivers as a high-risk group in terms of HTN. Besides, it was demonstrated that the variables of age, work history, WHPD, WDPW, sleep duration, economic status, and education level could be significantly associated with the prevalence of HTN among taxi drivers. More studies are required, especially longitudinal ones, to confirm the present study findings and to overcome the mentioned limitations.

Acknowledgement

The authors of the present study would like to appreciate the participants involved in this study.

Conflict of interest: None declared.

References

1. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *lancet* 2005; 365(9455):217-23.
2. Borzecki AM, Kader B, Berlowitz DR. The epidemiology and management of severe hypertension. *J Hum Hypertens* 2010; 24(1):9-18.
3. Lawes CM, Vander Hoorn S, Rodgers A; International Society of Hypertension. Global burden of blood-pressure-related disease, 2001. *Lancet* 2008; 371(9623):1513-8.
4. GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015; 385(9963):117-71.
5. GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018; 392(10159):1736-88.
6. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *lancet* 2012; 380(9859):2224-60.
7. GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018; 392(10159):1789-858.
8. Apantaku-Onayemi F, Baldyga W, Amuwo S, Adefuye A, Mason T, Mitchell R, et al. Driving to

- better health: cancer and cardiovascular risk assessment among taxi cab operators in Chicago. *J Health Care Poor Underserved* 2012; 23(2):768-80.
9. Burgel BJ, Gillen M, White MC. Health and safety strategies of urban taxi drivers. *J Urban Health* 2012; 89(4):717-22.
 10. Gany FM, Gill PP, Ahmed A, Acharya S, Leng J. "Every disease ... man can get can start in this cab": focus groups to identify south Asian taxi drivers' knowledge, attitudes and beliefs about cardiovascular disease and its risks. *J Immigr Minor Health* 2013; 15(5):986-92.
 11. Bruno PS, Marcos QR, Amanda C, Paulo ZH. Annoyance evaluation and the effect of noise on the health of bus drivers. *Noise Health* 2013; 15(66):301-6.
 12. Funakoshi M, Taoda K, Tsujimura H, Nishiyama K. Measurement of whole-body vibration in taxi drivers. *J Occup Health* 2004; 46(2):119-24.
 13. Brucker N, Charão MF, Moro AM, Ferrari P, Bubols G, Sauer E, et al. Atherosclerotic process in taxi drivers occupationally exposed to air pollution and co-morbidities. *Environ Res* 2014; 131:31-8.
 14. Wu S, Deng F, Niu J, Huang Q, Liu Y, Guo X. Association of heart rate variability in taxi drivers with marked changes in particulate air pollution in Beijing in 2008. *Environ Health Perspect* 2009; 118(1):87-91.
 15. Zagury E, Le Moullec Y, Momas I. Exposure of Paris taxi drivers to automobile air pollutants within their vehicles. *Occup Environ Med* 2000; 57(6):406-10.
 16. Hannerz H, Tüchsen F. Hospital admissions among male drivers in Denmark. *Occup Environ Med* 2001; 58(4):253-60.
 17. Bigert C, Gustavsson P, Hallqvist J, Hogstedt C, Lewné M, Plato N, et al. Myocardial infarction among professional drivers. *Epidemiology* 2003; 14(3):333-9.
 18. Tüchsen F, Hannerz H, Roepstorff C, Krause N. Stroke among male professional drivers in Denmark, 1994–2003. *Occup Environ Med* 2006; 63(7):456-60.
 19. Huang HY, Wang W, Zhou JP, Li QL, Feng WT, Wu ZZ. [Metabolic syndrome and its influencing factors in professional automobile drivers in a company]. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 2016; 34(4):258-61.
 20. Izadi N, Malek M, Aminian O, Saraei M. Medical risk factors of diabetes mellitus among professional drivers. *J Diabetes Metab Disord* 2013; 12(1):23.
 21. Mohebbi I, Saadat S, Aghassi M, Shekari M, Matinkhah M, Sehat S. Prevalence of metabolic syndrome in Iranian professional drivers: results from a population based study of 12,138 men. *PLoS One* 2012; 7(2):e31790.
 22. Siu SC, Wong KW, Lee KF, Lo YC, Wong CH, Chan AL, et al. Prevalence of undiagnosed diabetes mellitus and cardiovascular risk factors in Hong Kong professional drivers. *Diabetes Res Clin Pract* 2012; 96(1):60-7.
 23. Hege A, Lemke MK, Apostolopoulos Y, Sonmez S. Occupational health disparities among U.S. long-haul truck drivers: the influence of work organization and sleep on cardiovascular and metabolic disease risk. *PLoS One* 2018; 13(11):e0207322.
 24. Tabrizi R, Moosazadeh M, Razzaghi A, Akbari M, Heydari ST, Kavari SH, et al. Prevalence of sleep quality disorder among Iranian drivers: a systematic review and meta-analysis. *J Inj Violence Res* 2018; 10(1):53-9.
 25. Meng L, Zheng Y, Hui R. The relationship of sleep duration and insomnia to risk of hypertension incidence: a meta-analysis of prospective cohort studies. *Hypertens Res* 2013; 36(11):985-95.
 26. Li H, Ren Y, Wu Y, Zhao X. Correlation between sleep duration and hypertension: a dose-response meta-analysis. *J Hum Hypertens* 2019; 33(3):218-28.
 27. Liu Z, Wang Y, Yan F, Wei X, Yu S. Analysis of risk factors for hypertension among taxi drivers on different shifts. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 2015; 33(4):263-5.
 28. Shin SY, Lee CG, Song HS, Kim SH, Lee HS, Jung MS, et al. Cardiovascular disease risk of bus drivers in a city of Korea. *Ann Occup Environ Med* 2013; 25(1):34.
 29. Elshatarat RA, Burgel BJ. Cardiovascular Risk Factors of Taxi Drivers. *J Urban Health* 2016; 93(3):589-606.
 30. Adedokun AO, Ter Goon D, Owolabi EO, Adeniyi OV, Ajayi AI. Driving to Better Health: Screening for Hypertension and Associated Factors among Commercial Taxi Drivers in Buffalo City Metropolitan Municipality, South Africa. *Open Public Health J* 2017; 10(1):303-12.
 31. Persu A, Andries A, Demedts S, Van der Niepen P, van de Borne P, Belgian Hypertension Committee. Elevated prevalence of arterial hypertension amongst Belgian taxi drivers during the World Hypertension Day campaign 2006. *J Hypertens* 2006; 24(11):2311-12.
 32. Cavagioni LC, Pierin AM, Batista KM, Bianchi ER, Costa AL. Health problems, arterial hypertension and predisposition to stress in truck drivers. *USP School of Nursing Magazine* 2009; 43(2):1267-71.
 33. Lakshman A, Manikath N, Rahim A, Anilakumari VP. Prevalence and Risk Factors of Hypertension among Male Occupational Bus Drivers in North Kerala, South India: A Cross-Sectional Study. *ISRN Prev Med* 2014; 2014:318532.
 34. Tobin EA, Ofili AN, Asogun DA, Igbinosun PO, Igba K, Idahosa A. Prevalence of hypertension and associated factors among inter-city drivers in an urban city in South-South Nigeria. *Int J Res Med* 2013; 2(3):5-12.

35. Oyeniyi OS, Ajayi IO. Prevalence of hypertension and associated risk factor among interstate commercial drivers in Jabi Park Abuja. *International Journal of Medicine and Medical Sciences* 2016; 8(7):75-83.
36. Platek AE, Szymanski FM, Filipiak KJ, Kotkowski M, Rys A, Semczuk-Kaczmarek K, et al. Prevalence of Hypertension in Professional Drivers (from the RACER-ABPM Study). *Am J Cardiol* 2017; 120(10):1792-6.
37. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, et al. 2018 ESC/ESH Guidelines for the management of arterial hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology (ESC) and the European Society of Hypertension (ESH). *Eur Heart J* 2018; 39(33):3021-104.
38. Gany F, Bari S, Gill P, Ramirez J, Ayash C, Loeb R, et al. Step On It! Workplace Cardiovascular Risk Assessment of New York City Yellow Taxi Drivers. *J Immigr Minor Health* 2016; 18(1):118-34.
39. Martin WP, Sharif F, Flaherty G. Lifestyle risk factors for cardiovascular disease and diabetic risk in a sedentary occupational group: the Galway taxi driver study. *Ir J Med Sci* 2016; 185(2):403-12.
40. Yang Y, Fan XS, Tian CH, Zhang W, Li J, Li SQ. Health status, intention to seek health examination, and participation in health education among taxi drivers in Jinan, China. *Iran Red Crescent Med J* 2014; 16(4):e13355.
41. Vieira MC, Sperandei S, Reis AC. Physical activity overcomes the effects of cumulative work time on hypertension prevalence among Brazilian taxi drivers. *J Sports Med Phys Fitness* 2016; 56(5):631-8.
42. Khoshandam Sarvynnebaghi F, Mozaffari SAR, Yaghoubi Poor A, Nezamtabar Malekshah A. Prevalence of Hypertension among Professional Drivers in Mazandaran Province, 2010. *Journal of Mazandaran University of Medical Sciences* 2013; 23(102):19-24.
43. Borle AL, Jadhao A. Prevalence and associated factors of hypertension among occupational bus drivers in Nagpur City, Central India-A cross sectional study. *Age (Years)* 2015; 6:423-8.
44. Hirata RP, Sampaio LM, Leitão Filho FS, Braghiroli A, Balbi B, Romano S, et al. General characteristics and risk factors of cardiovascular disease among interstate bus drivers. *ScientificWorldJournal* 2012; 2012:216702.
45. dos Reis LAP, Costa CDD, Rodrigues DS, de Alcântara KC. Obesity, hypertension and diabetes among truck drivers in the middle-west, Brazil. *Biosci J* 2017; 33(2):485-93.
46. Saberi HR, Moravveji AR, Fakharian E, Kashani MM, Dehdashti AR. Prevalence of metabolic syndrome in bus and truck drivers in Kashan, Iran. *Diabetol Metab Syndr* 2011; 3(1):8.
47. Buford TW. Hypertension and aging. *Ageing Res Rev* 2016; 26:96-111.
48. Bawa MS, Srivastav M. Study the epidemiological profile of taxi drivers in the background of occupational environment, stress and personality characteristics. *Indian J Occup Environ Med* 2013; 17(3):108-13.
49. Leng B, Jin Y, Li G, Chen L, Jin N. Socioeconomic status and hypertension: a meta-analysis. *J Hypertens* 2015; 33(2):221-9.
50. Satheesh B, Veena RM. A study of prevalence of hypertension among bus drivers in Bangalore City. *Int J Curr Res Rev* 2013; 5:90-4.