



Natural Sounds in Management of Elderly Sleep Quality: A Randomized Controlled Trial

Ali Ansari Jaber¹, Zohreh Sahebi², Tayebeh Negahban Bonabi^{3*}, Poursan Allah Bakhshinasab⁴

1. Instructor, Dept. of Psychiatric and Mental Health Nursing, Social Determinants of Health Research Center, School of Nursing and Midwifery; Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

2. Instructor, Dept. of Midwifery, School of Nursing and Midwifery, Geriatric Care Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

3. Assistant Prof., Dept. of Community Health Nursing, Social Determinants of Health Research Center, School of Nursing and Midwifery, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

4. Instructor, Dept. of Midwifery, School of Nursing and Midwifery Geriatric Care Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.



Citation: Ansari Jaber A, Sahebi Z, Negahban Bonabi T, Allah Bakhshinasab P. Natural Sounds in Management of Elderly Sleep Quality: A Randomized Controlled Trial. *J Occup Health Epidemiol.* 2023;12(4):213-9.

Copyright: © 2023 The Author(s); Published by Rafsanjan University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

Article Info

* **Corresponding author:**
Tayebeh Negahban Bonabi,
E-mail:
negahbant@yahoo.com

Article history

Received: Jan 2023

Accepted: Oct 2023

doi 10.61186/johe.12.4.213

Print ISSN: 2251-8096

Online ISSN: 2252-0902

Peer review under responsibility of
Journal of Occupational Health
and Epidemiology

Abstract

Background: There is no clear consensus on the effect of natural sounds on elderly sleep quality, as one of the most common causes of disability in them. This study aimed to determine the effect of natural sounds on elderly sleep quality.

Material & Methods: In this randomized controlled trial, 100 elderly were selected based on the inclusion criteria from comprehensive health care centers and then divided into two equal groups by simple random sampling method. The intervention group listened to the natural sounds in MP3 format every night for 2 weeks, 30 minutes before going to bed, through mobile phone and hands-free. The control group did not receive any intervention. The PSQI was measured before and after the intervention. Data analysis was done in SPSS software (version 22) applying the Shapiro-Wilk test, Chi-square, and Fisher Exact test, Independent Samples-t test, Wilcoxon Signed Ranks test, and Mann Whitney U test. A confidence level of 95% was considered.

Results: In the intergroup comparison, in the baseline, groups were the same in terms of the total sleep quality score as well as other components ($P > 0.05$). After the intervention, there was a significance difference between the two groups in terms of global sleep quality ($P = 0.048$), sleep latency ($P = 0.001$), habitual sleep efficacy ($p = 0.012$), and sleep disturbance ($p = 0.01$).

Conclusion: Based on the our results, natural sounds could improve the sleep quality in the elderly.

Keywords: Sleep Qualities, Elderly, Sounds.

Introduction

Along with physical changes, changes in sleep patterns are also part of the natural aging process [1]. About 43% of older persons report having trouble falling asleep and staying asleep as they become older. Older adults also often have early wake times and sleeplessness. They spend the majority of their sleep in

light sleep phases. [2]. Researchers found that the prevalence of poor sleep quality was 33.8% in elderly people[3].

Good sleep is necessary for health. Sleep disorders in the elderly are associated with serious negative physical, psychological, and social consequences(4).A recent research shows that insufficient sleep, sleep disturbance, and sleep disorders affect many aspects of human

health, including cognitive decline, Alzheimer's disease risk(5-7) dementia(8) polypharmacy (1), sexual performance [9], and consequently the quality of life [10, 11] and morbidity and mortality rate (1). Also the results of studies showed that other common medical problems in old age, such as high blood pressure, diabetes mellitus, kidney failure, immune system disorders, gastroesophageal reflux disease, respiratory diseases such as asthma, physical disability, depression, and anxiety and dementia are all associated with sleep disorders(12).

Researchers proposed several approaches to deal with sleep disorders in the elderly, such as physiological and pharmacological methods(4). Some of the non-pharmacological interventions are: natural light therapy(13, 14), brain exercise(15) cognitive behavioral(16) physical activity(17) aromatherapy(18) medicinal Plants(19), auditory stimulation(20) like several types of music therapy (21), include analytical music therapy, cognitive behavioral music therapy (CBMT), community music therapy, Benenson music therapy, vocal psychotherapy and Nordoff-Robbins music therapy, (22), and probably natural sounds(23).

Researchers have recently demonstrated that the sounds of nature induce a state of relaxation by modulating the central and autonomic nervous systems (24). Additionally, some participants have reported that listening to music prior to napping enhances both the subjective and objective parameters of their sleep (21, 25, 26). The effect of natural sounds on sleep quality have been studied in limited studies, and most of the available reports are based on the results of studies conducted on the effect of music on sleep quality in hospitalized patients. Limited studies showed that, compared to silence, nature sounds did not show significant effects on patients quality of sleep(27). Regarding conflicting reports about the effects of music therapy on sleep quality, researchers in the results of a systematic review, have recommended more studies in this field(28). This study aimed to determine the effect of natural sounds on elderly sleep quality.

Materials and Methods

The design of this study was a randomized controlled trial performed from May 2022 to August 2022; to determine the effect of natural sounds on elderly sleep quality, Rafsanjan, Iran.

Research Council of Rafsanjan University of Medical Sciences approved the project. The code of ethics (IR.RUMS.REC.1399.133) was adopted by the ethics committee by deputy of the research, and technology at this university, the permission to conduct the study was present in the comprehensive health service centers of Rafsanjan by two of the researchers. With this end in view, first, they prepared a list of elderly people (85

people) who had a history of sleep disorders at health electronic files at the Iranian Health Integrated System (SIB), and then through a phone call. The researchers invited the elderly who satisfied the inclusion criteria and consented to participate in the study to the comprehensive health care facilities after outlining the goal of the investigation. When invited to comprehensive health care facilities, almost all of the individuals consented to take part in the study. The inclusion criteria include age over 60 years(29), no experiencing a life crisis during the last 6 weeks, obtaining a total score higher than 5 from Pittsburgh Sleep Quality Index (PSQI), no hearing problems, having sufficient mental and cognitive ability to answer the questionnaire and to use hands-free, or the presence of family members who can help them play audio through hands-free, no addiction such as alcohol, opium, smoking or tranquilizers and sleeping drugs.

Exclusion criteria included: forgetting more than three days to implement the intervention, withdrawing from continuing to attend the study, and occurrence of any problem that implements the intervention impossible.

The sample size considering $\alpha \leq 0.05$, the $\beta = 0.90$ and $S_1 = 2.91$ and $S_2 = 3.03$ for control and intervention groups respectively, and $d = 3$ for the 1 hour per day for 1 week intervention(30) and using the following equation is approximately 41 people calculated. But we considered 50 people for each group.

Formula 1.

$$n = 2(z_{1-\alpha} + z_{1-\beta})^2 (S_1^2 + S_2^2) / d^2$$

At the beginning of study, the samples were selected purposefully, and based on the inclusion criteria, using the SIB system then the eligible samples were then allocated into two groups of the intervention (natural sound) and control (without intervention), by a simple random lottery method.

The data collection tool consists of two parts. The first was the demographic characteristics (including age, sex, level of education, job, income, and sleep disorder duration), and the second part was the PSQI. PSQI is a standard scale that has 19 items and 7 subscales (sleep quality, sleep delay, and sleep duration, normal sleep efficiency, sleep disorders, using sleeping pills, and daily functioning disorders). Each component is scored from 0 to 3 (not in the past month = 0; less than once a week = 1, once or twice a week = 2, and three or more times a week = 3). The minimum is 0 and the maximum is 21 variables. Higher scores indicate lower sleep quality. The total sleep score of 5 or less is considered "good"(31). In the Persian version the Cronbach's Alpha Coefficient of 0.77 was reported for all subjects. The psychometric properties of the Persian version of the PSQI were approved (32).

To reduce the interfering effects of environmental factors on sleep, before going to sleep, the elderly of both groups were advised to be in a calm, quiet, low-light environment with a suitable temperature, where, they feel comfortable, wear light, and comfortable clothes, lie down in their bed and a comfortable position. The samples were advised to observe the following instructions, before listening to the audio file. They should close their eyes and concentrate on the sound. Then, while maintaining regular respiration, they should adjust the volume of the sound to a level that is suitable for them. They should not be concerned with turning off their mobile phone. Except for preliminary recommendations at the beginning of sleep, no other intervention was implemented for the control group.

The intervention group, listened to the natural sounds in MP4 format every night for 2 weeks, 30 minutes before going to sleep, through mobile phone and hands-free. The nature sounds in this study were the artwork "Sounds of Nature" which was recorded by Dr. Arendt Stein in 2011. These sounds included the chirping of birds, the roar of the sea, the sounds of river and the forest, and the sound of rain. It was recommended that the volume of sounds be adjusted to a level that, feels comfortable.

For all participants in this study, PSQI was measured the day before the start of the intervention, and the morning after the last night of the intervention. The baseline measurement was done through a face-to-face interview and the second time measurement was done at least 8 hours after the last intervention (at 10:00 am), through a phone call interview. Both measurements were done by the researcher's colleague who was blind about the sample allocation.

Ethical considerations were upheld, including the following: voluntary participation and termination of the study at will; acquisition of informed written consent prior to study commencement; and protection of information confidentiality. The samples were assured that the intervention is not associated with any physical, psychological, spiritual, or financial complications to them.

Data were analyzed by SPSS software version 22 by using a descriptive statistic, Shapiro -Wilk test (for normality) Chi-square and Fisher Exact test (to compare ratios), Independent samples-t test, and Wilcoxon Signed Ranks test (For intragroup comparisons) Mann Whitney U test (For intergroup comparisons). A confidence level of 95% was considered

Results

Based on Shapiro-Wilk test, except for the age variable, the quantitative data distribution was not normal. Data analysis results showed that, in terms of gender, 52 (52%) of the samples were male and 42 (42%) were female, the mean ± SD of the elderly age was 68.66 ± 5.37 with a maximum of 82 and a minimum of 60 years. The independent samples-t test showed that the mean ± SD of age in intervention and control groups were respectively 68.06± 5.66 and 69.26±5.046. Regarding age, no statistically significant difference existed between the groups under investigation (p=0.266). Additional demographic factors, including sleep disorder duration, occupation, gender, and occupation, did not exhibit any statistically significant variations among the study groups. Table 1 shows the study groups' similarity in terms of these features.

Table 1. Comparison of demographic characteristics between the study groups

		Intervention n=50 (100%)	Control n=50 (100%)	Pearson chi-square test P-value
Gender	Male	26(52)	32(64)	0.224*
	Female	24(48)	18(36)	
Education	Illiterate	21(42)	20(40)	0.085*
	Diploma & under diploma	11 (22)	20(40)	
	Academic	18(36)	10 (20)	
Job	Unemployed	21(42)	20(40)	0.500**
	Retired	28(56)	26(52)	
	Employed	1(2)	4(8)	
History of sleep disorder	Less than one month	9(18)	16(32)	0.106*
	More than one month	41(82)	34(68)	

* Chi-square test
**Fisher Exact test

The mean and standard deviation of total sleep score of the intervention group, in the pre-intervention phase was 10.24±3.47 and reached in the post-intervention phase to 8.44±3.86. In this group, in within-group comparison, a significant reduction in global quality of sleep score

(p=0.001), subjective sleep score (p=0.001), and sleep latency (p=0.001) showed in results of the Wilcoxon Signed Rank test. In other components of sleep quality, no statistically significant difference was observed (P>0.05) (Table 2).

Table 2. The intra-group comparison of global quality of sleep score and other components in the intervention group

	Pre-test Mean±SD	Post-test Mean±SD	Pre-test Median±IQR	Post-test Median±IQR	*P-value
Global sleep score	10.24±3.47	8.44±3.86	5.25±9	7.50±5.25	0.001
Subjective sleep quality	1.68±0.84	1.20±0.94	2±1	1±1	0.001
Sleep latency	2.44±0.61	1.70±0.83	2.50±1	2±1	0.001
Sleep duration	2.10±0.99	1.90±1.01	2±1	2±2	0.057
Habitual sleep efficacy	0.8±0.92	0.72±1.05	1±1	0±1.25	0.432
Sleep disturbance	1.52±0.58	1.40±0.53	1±1	1±1	0.058
Use of sleep medication	0.76±1.20	0.72±1.17	0±1	0±1	0.317
Daytime dysfunction	0.96±0.85	0.84±0.88	1±2	1±2	0.058

*Wilcoxon Signed the Rank test

The mean ± SD of the global quality of sleep score of the control group in the pre-intervention phase was 10.18±3.70 and in the post-intervention phase reached 9.88±3.57. Results of the Wilcoxon Signed Rank test showed that, except for the sleep latency, there was no significant difference between the global quality of sleep score and scores of the other components in the pre-and post-measurement phases (P>0.05). In the control group, the mean ± SD of the sleep latency in the pre-intervention phase was 2.46±0.73 and reached in the post-intervention phase to 2.26±0.92, which was statistically significance (P=0.029).

In the intergroup comparison, in the pre-intervention phase, both groups were similar in terms of the global quality of sleep score and other subscales (P>0.05). In post post-intervention phase, there was a statistical significance difference between study groups in terms of PSQI score (P=0.048), sleep latency (P=0.001), habitual sleep efficacy (p= 0.012), and sleep disturbance (p=0.01). Other components' scores were the same between the two groups (P>0.05) (Table 3). Due to the use of non-parametric tests, comparisons with median and quartile ranges are also reported in this table.

Table 3. The inter-group comparison of global quality of sleep score and other components across study groups

		Intervention group Mean±SD	Control group Mean±SD	Intervention group Median±IQR	Control group Median± IQR	*P-value
Global sleep score	Pre-test	10.24±3.47	10.18±3.70	9±5.25	10.00±	0.805
	Post-test	8.44±3.86	9.88±3.57	7.25±5.25	9.50±5.25	0.048
Subjective sleep quality	Pre-test	1.68±0.84	1.48±0.73	2±1	1±1	0.231
	Post-test	1.20±0.94	1.38±0.63	1±1	1±1	0.124
Sleep latency	Pre-test	2.44±0.61	2.46±0.73	2.50±1	3±1	0.600
	Post-test	1.70±0.83	2.26±0.92	2±1	3±1	0.001
Sleep duration	Pre-test	2.10±0.99	1.92±1.00	2±2	2±2	0.345
	Post-test	1.90±1.01	2.00±0.85	2±2	2±2	0.685
Habitual sleep efficacy	Pre-test	0.8±0.92	0.92±0.85	1±1	1±1	0.325
	Post-test	0.72±1.05	1.04±0.83	0±1.25	1±0.25	0.012
Sleep disturbance	Pre-test	1.52±0.58	1.72±0.57	1±1	2±1	0.075
	Post-test	1.40±0.53	1.72±0.62	1±1	2±1	0.010
Use of sleep medication	Pre-test	0.76±1.20	0.74±1.08	0±1	0±1.25	0.869
	Post-test	0.72±1.17	0.74±1.06	0±1	0±2	0.724
Daytime dysfunction	Pre-test	0.96±0.85	0.96±1.02	1±2	1±1	0.704
	Post-test	0.84±0.88	0.86±0.92	1±2	1±1	0.971

* Mann-Whitney U test

Discussion

The results of the study stated that, in the intervention group, the natural sounds were able to improve significantly the sleep quality score compared to the baseline, and compared to the control group. This results were similar to study by Afshar et al., on sleep quality management by white noise in CCU patients, on the third day after admission. Results of this study showed that white noise in the intervention group, could

improve the quality of sleep, compared to the before intervention and the control group(33). Similar to the work of Afshar and colleagues, Nasari et al., conducted a study in the coronary care patients, by the natural sounds, and adding a third group (silence group). The researchers reached the conclusion that the quality of sleep was enhanced in both the silence and natural sounds intervention groups in comparison to the control group. However, no significant difference was observed

between the silence and natural sounds group. These researchers concluded that based on the patients' preference, silence or nature sounds can be used to improve the quality of sleep of the patients(27). In other studies, person-preferred music therapy was introduced as a painless, safe, affordable treatment method without any side effects intervention to improve sleep quality(34-36). Also, the results of Lan and colleagues' study showed that the speed of ventilation, and noise of ventilation are the influencing factors on the quality of sleep of patients hospitalized in ICU(37). Undoubtedly, the environment of the intensive care units, and the contextual causes of sleep quality reduction in patients hospitalized in these units are different from the elderly own living environments. However, it should be kept in mind that, most of the patients hospitalized in these units are elderly. The results of these studies showed that environmental noises were an important factor to reduce the sleep quality of patients who were hospitalized in intensive care units, so that, the elimination of these sounds in the silence group could improve the sleep quality of the patients as well as in the natural sounds group. Although the problem of these patients can be different in nature from the elderly who have suffered a long-term loss of sleep quality in their real-life environment. Therefore, finding the appropriate strategies for the management of their sleep quality can be critical. Based on the results of this study, besides the adjusting of environmental factors (such as temperature, light, and noise), at home and in a personal and regular bed, the natural sounds could improve the sleep quality of the elderly. The positive impact of natural music on sleep quality can be ascribed to its calming and tranquilizing properties, which serve to divert attention away from negative thoughts. A study's findings indicate that music therapy has the capacity to decrease anxiety, pulse, and respiration rate, while also maintaining systolic and diastolic blood pressure within the expected range (38). Apart from the score of sleep quality, although the sleep latency score (the duration of falling asleep) was improved in both groups at post-intervention, but in the music therapy group, the sleep latency score was shorter and better than the control group. Moreover, the efficiency of sleep (times of going to bed and morning waking up and actual sleep duration habits), and sleep disturbance (waking up in the middle of the night or early in the morning) were significantly improved in the intervention group. The results of some other studies, including a systematic review meta-analysis study showed continuous improvements in sleep quality, in global PSQI score compared to the baseline, and three consecutive months after the music intervention, also, the score of sleep quality in the intervention group compared to the control group had improved more in each measurement time. Three

subscale scores including sleep efficiency, sleep latency, and daytime dysfunction were improved in intervention groups (21, 39, 40).

Scientists believe that the sounds of nature, inspired by music and poetry, create powerful reactions in humans and connect them to nature by creating a sense of place (41). Hence, it can improve the health and well-being of humans, increase positive affect, and reduce stress and anxiety(42), all of which can be beneficial in providing healthy sleep and improving sleep quality.

Despite trying to design of the study to obtain valid results by eliminating environmental factors, removing anthropogenic sounds in order to provide pure biological sounds, and maintaining blinding in data collection, but, some limitations of this study, such as the effect of frequent therapist calls on the feeling of attention, security, and peace of the elderly in the intervention group and how seniors answer skill questions. Thus, the effect of the patient's lifestyle such as activity level, feeding style, daily events, and their housemate's conditions that, could affect on their quality of sleep was beyond our precise control. Further investigation is recommended in order to provide a rationale for the restricted efficacy of interventions such as nature sounds and music therapy with regard to other facets of sleep. Furthermore, further research is warranted to ascertain the precise populations and contexts in which this approach may prove to be therapeutically beneficial.

Conclusion

Based on the results of this study, natural sounds improve sleep quality in the elderly. This effect is the result of changes in the fields of sleep latency score, also, the efficiency of sleep, and sleep disturbance. But, other aspects of sleep did not change. Nurses and other health care professionals can implement these interventions as a possible effective method for the quality of sleep, taking into account cultural contexts for the elderly, to improve their quality of sleep.

Acknowledgement

This study is extracted from the master's thesis. Unfortunately, Mr. Maitham Afzali, the relevant student, lost his life while caring for patients suffering from Covid-19 and he was unable to complete his work. While thanking, we hereby appreciate his efforts and honor his memory. Furthermore, the authors would like to express their gratitude to all participants in this study and the Deputy of Research and Technology of Rafsanjan University of Medical Sciences for their financial support.

Conflict of interest: None declared.

References

1. Miner B, Kryger MH. Sleep in the Aging Population. *Sleep Med Clin*. 2017;12(1):31-8.
2. Patel AK, Reddy V, Shumway KR, Araujo JF. Physiology, sleep stages. In: StatPearls [Internet]: StatPearls Publishing; 2022. Available from: https://www.ncbi.nlm.nih.gov/books/NBK526132/#_NBK526132_dtls__
3. Wang P, Song L, Wang K, Han X, Cong L, Wang Y, et al. Prevalence and associated factors of poor sleep quality among Chinese older adults living in a rural area: a population-based study. *Aging Clin Exp Res*. 2020;32(1):125-31.
4. Rodriguez JC, Dzierzewski JM, Alessi CA. Sleep problems in the elderly. *Med Clin North Am*. 2015;99(2):431-9.
5. Irwin MR, Vitiello MV. Implications of sleep disturbance and inflammation for Alzheimer's disease dementia. *Lancet Neurol*. 2019;18(3):296-306.
6. You JC, Jones E, Cross DE, Lyon AC, Kang H, Newberg AB, et al. Association of β -amyloid burden with sleep dysfunction and cognitive impairment in elderly individuals with cognitive disorders. *JAMA Netw Open*. 2019;2(10):e1913383.
7. Brzecka A, Leszek J, Ashraf GM, Ejma M, Ávila-Rodríguez MF, Yarla NS, et al. Sleep disorders associated with Alzheimer's disease: a perspective. *Front Neurosci*. 2018;12:330.
8. Shi L, Chen SJ, Ma MY, Bao YP, Han Y, Wang YM, et al. Sleep disturbances increase the risk of dementia: a systematic review and meta-analysis. *Sleep Med Rev*. 2018;40:4-16.
9. Cho JW, Duffy JF. Sleep, sleep disorders, and sexual dysfunction. *World J Mens Health*. 2019;37(3):261-75.
10. Uchmanowicz I, Markiewicz K, Uchmanowicz B, Kołtuniuk A, Rosińczuk J. The relationship between sleep disturbances and quality of life in elderly patients with hypertension. *Clin Interv Aging*. 2019;14:155-65.
11. Lubetkin EI, Jia H. Burden of disease due to sleep duration and sleep problems in the elderly. *Sleep Health*. 2018;4(2):182-7.
12. Gulia KK, Kumar VM. Sleep disorders in the elderly: a growing challenge. *Psychogeriatrics*. 2018;18(3):155-65.
13. Fery Agusman Motuho Mendrofa F, Aeni M. Natural light therapy to lower agitation and sleep disturbance of dementia patients in semarang. *Int J Psychosoc Rehabil*. 2020;24(7):7752-9.
14. Abidin Z, Yasmara D. The Effectiveness of Afternoon Sunbath in Improving Elderly Sleep Quality. *Indian J Public Health Res Dev*. 2019;10(9).
15. Irvana I, Alvita GW, Pramudaningsih IN, Fitriana V. Therapy of Brain Exercise on The Quality of Sleeping in Elderly. *South East Asia Nurs Res*. 2020;2(3):88-93.
16. Chokroverty S. Sleep disorders in the elderly. In: Chokroverty, S. (eds). *Sleep disorders medicine*. New York, NY, United States: Springer; 2017. P.1115-38.
17. Podhorecka M, Cytarska M, Gębka D, Perkowski R, Androsiuk-Perkowska J, Jaroch A, et al. Can physical activity influence the quality of sleep among the elderly? *J Educ Health Sport*. 2017;7(12):288-305.
18. Wasijati W, Waluyo A, Maria R. The Effect of Aromatherapy Massage on Sleep Quality of Patients with Cancer. *Proceedings of the Third International Conference on Sustainable Innovation 2019–Health Science and Nursing (IcoSIHSN 2019)*; 2019 Oct; *Adv Health Sci Res*. 2019;15:157-61. Atlantis Press.
19. Subi Mol R. An experimental study to assess the effectiveness of Chamomile Tea on Insomnia and Anxiety among Institutionalized elderly in selected old age homes, Thanjavur. [Phd thesis]. Thanjavur, India: Our Lady of Health College of Nursing; 2016.
20. Capezuti E, Pain K, Alamag E, Chen X, Philibert V, Krieger AC. Systematic review: auditory stimulation and sleep. *J Clin Sleep Med*. 2022;18(6):1697-709.
21. Wang C, Li G, Zheng L, Meng X, Meng Q, Wang S, et al. Effects of music intervention on sleep quality of older adults: A systematic review and meta-analysis. *Complement Ther Med*. 2021;59:102719.
22. Law S, Au DC, Leung AW, Xu C. Music therapy for elderly sleeping quality. *Geriatr Care*. 2022;8(1).
23. Yoon H, Baek HJ. External Auditory Stimulation as a Non-Pharmacological Sleep Aid. *Sensors (Basel)*. 2022;22(3):1264.
24. Hikaru H, Keijiro S, Okada S, Hiromatsu K, Takeda N. The Relaxation Effect of Music and Natural Sounds. Paper presented at: The 4th Global Conference on Life Sciences and Technologies (LifeTech); 2022 Mar 7-9; Osaka, Japan.
25. Altan Sarikaya N, Oğuz S. Effect of Passive Music Therapy on Sleep Quality among Elderly People Living in Nursing Home. *J Psychiatr Nure*. 2016;7(2):55-60.
26. Lestarini A, Putra KSP. Music therapy on sleep quality in elderly. *IOP Conf Ser Mater Sci Eng*. 2018;434.012143.
27. Nasari M, Najafi Ghezeljeh T, Haghani H. Effects of nature sounds on sleep quality among patients hospitalized in coronary care units: A randomized controlled clinical trial. *Nurs Midwifery Stud*. 2018;7(1):18-23.
28. Pan BY. The use of music therapy/music to address sleep quality and sleep disorders: a systematic literature review. [MSc Thesis]. Quebec, Canada: Concordia University; 2017.
29. Bahramnezhad F, Farokhnezhad Afshar P, Asgari P, Shiri M. A review of the Iran's elderly status according to the census records. *Galen Med J*. 2016;5(1):e397.
30. Najafi Ghezeljeh T, Nasari M. Investigating the effect of listening to nature sounds on sleep quality in patients hospitalized in cardiac care units. *Complement Med J*. 2018;8(1):2167-80.
31. Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193-213.
32. Farrahi Moghaddam J, Nakhaee N, Sheibani V, Garrusi B, Amirkafi A. Reliability and validity of the

- Persian version of the Pittsburgh Sleep Quality Index (PSQI-P). *Sleep Breath.* 2012;16(1):79-82.
33. Farokhnezhad Afshar P, Bahramnezhad F, Asgari P, Shiri M. Effect of white noise on sleep in patients admitted to a coronary care. *J Caring Sci.* 2016;5(2):103-9.
34. Kavurmacı M, Dayapoğlu N, Tan M. Effect of music therapy on sleep quality. *Altern Ther Health Med.* 2020;26(4):22-6.
35. Loewy J. Music therapy as a potential intervention for sleep improvement. *Nat Sci Sleep.* 2020;12:1-9.
36. Huda KK. Natural Sound Music Therapy on Sleep Disorders of Post Operation Patients. *J Appl Health Manag Technol.* 2022;4(3):42-5.
37. Lan L, Sun Y, Wyon DP, Wargocki P. Pilot study of the effects of ventilation and ventilation noise on sleep quality in the young and elderly. *Indoor Air.* 2021;31(6):2226-38.
38. Pilang A, Dyah DI, Pujiastuti RS. Effect of "Sape" Music Therapy on Decreasing Anxiety, Physiological Response, and Sleep Quality among Post Sectio Caesarea Patients. *Int J Nurs Health Serv.* 2022;5(1):28-37.
39. Wang Q, Chair SY, Wong EM, Li X. The effects of music intervention on sleep quality in community-dwelling elderly. *J Altern Complement Med.* 2016;22(7):576-84.
40. Feng F, Zhang Y, Hou J, Cai J, Jiang Q, Li X, et al. Can music improve sleep quality in adults with primary insomnia? A systematic review and network meta-analysis. *Int J Nurs Stud.* 2018;77:189-96.
41. Erfanian M, Mitchell AJ, Kang J, Aletta F. The psychophysiological implications of soundscape: A systematic review of empirical literature and a research agenda. *Int J Environ Res Public Health.* 2019;16(19):3533.
42. Buxton RT, Pearson AL, Allou C, Fristrup K, Wittemyer G. A synthesis of health benefits of natural sounds and their distribution in national parks. *Proc Natl Acad Sci U S A.* 2021;118(14):e2013097118.