




Effects of Biomechanical Risk Factors as Cumulative Trauma on Rotator Cuff Tendinopathy in Workers - A Clinical Survey (2018 - 2019)

Saber Mohammadi¹, Fatemeh Shidfar², Seyed Reza Saadat Mostafavi³, Reza Salehi⁴, Afshin Zarafshar⁵, Elaheh Kabir-Mokamelkhan^{6*}

1. Professor, Dept. of Occupational Medicine, School of Medicine, Iran University of Medical Sciences, Tehran, Iran.
2. MD, Dept. of Occupational Medicine, School of Medicine, Iran University of Medical Sciences, Tehran, Iran.
3. Assistant Prof., Dept. of Radiology, School of Medicine, Hazrat-e Rasool General Hospital, Iran University of Medical Sciences, Tehran, Iran.
4. MD, Dept. of Radiology, Fayazbaksh Hospital, Tehran, Iran.
5. MD, Dept. Occupational Medicine, School of Medicine, Iran University of Medical Sciences, Tehran, Iran.
6. Associated Prof., Occupational Medicine Research Center (OMRC), Dept. of Occupational Medicine, School of Medicine, Iran University of Medical Sciences, Tehran, Iran.


 **Citation:** Mohammadi S, Shidfar F, Saadat Mostafavi SR, Salehi R, Zarafshar A, Kabir-Mokamelkhan E. Effects of Biomechanical Risk Factors as Cumulative Trauma on Rotator Cuff Tendinopathy in Workers - A Clinical Survey (2018 - 2019). *J Occup Health Epidemiol.* 2022;11(3):223-30.

Article Info

* **Corresponding author:**
Elaheh Kabir-Mokamelkhan,
E-mail:
kabir.e@iums.ac.ir

Article history

Received: Jan 2022
Accepted: Aug 2022

 10.52547/johe.11.3.223

Print ISSN: 2251-8096
Online ISSN: 2252-0902

Peer review under
responsibility of Journal of
Occupational Health and
Epidemiology

Abstract

Background: Cumulative effects of biomechanical risk factors on rotator cuff tendinopathy have not yet been determined. Accordingly, this study aimed to investigate cumulative effects of biomechanical risk factors on rotator cuff tendinopathy among workers.

Materials & Methods: This descriptive study was conducted on 470 patients suspected of rotator cuff tendinopathy, who referred to the radiology department of Rasool-e-Akram Hospital in Tehran, Iran (July 2018-March 2019). To this end, ergonomic risk factors, including excessive hand raising, repetitive tasks, upper arm flexion, grip force, awkward posture, vibration, abduction, and rotation were evaluated. The subjects were divided into four categories, according to the number of affecting risk factors. Next, univariate and multivariate analyses were performed to identify predictors of rotator cuff tendinopathy. The study data were analyzed by SPSS V24.0.

Results: Risk factors, such as excessive hand raising, repetitive tasks, upper arm flexion, and awkward posture had a significant association with the frequency of rotator cuff tendinopathy ($p < 0.05$). Our study showed that shoulder tendinopathy had a statistically significant association with having at least one affecting risk factor (OR=3.96 95%CI: 2.26-6.96), two affecting risk factors (OR=6.82 95%CI: 4.13-11.26), three affecting risk factors (OR=10.25 95%CI: 4.35-24.14), and four affecting risk factors (OR=12.61 95%CI: 1.69-93.63).

Conclusion: The existence of more than one mechanical risk factor in the workplace is associated with a greater increase in the possibility of rotator cuff tendinopathy. In most workplaces, there is usually more than one mechanical risk factor, so it is necessary to pay attention to cumulative effects of all of them.

Keywords: Shoulder Pain, Posture, Risk Factors, Workplace, Workers.

Introduction

According to research, the lifetime prevalence of shoulder pain in the general population and

patients less than 70 years old was 67 and 27%, respectively [1]. In a large-scale population-based survey in Australia in 2010, the prevalence of

shoulder pain was reported to be 22% [2]. Shoulder pain can cause sickness absence, loss of personal ability, as well as reduced work performance and work efficiency [3]. One of the common causes of shoulder pain and upper extremity musculoskeletal disorders (MSDs) in the general population is structure tendinopathy that includes structure tendinitis/tendinosis, subacromial bursitis, and impingement syndrome [4-13]. Shoulder pain improves slowly over time, so the likelihood of recurrence is comparatively high, with about half (40-50%) of patients with shoulder pain suffering from sustained pain or recurrence during the 12-month follow-up period [14, 15]. Successful treatment of structure tendinopathy remains challenging and is especially conservative, with about 40% of patients with structure tendinopathy not responding to such treatments [16-18]. Thus, it is necessary to identify the risk factors and causes of structure tendinopathy for the purpose of prevention interventions. The etiology of structure tendinopathy is multifactorial, so both intrinsic and extrinsic factors are involved [19]. The major mechanism of structure tendinopathy is the narrowing of the subacromial space and repetitive overhead activities, such as tendon overuse and overload [20]. Several studies show that diabetes mellitus and abdominal obesity are considered intrinsic factors, with work-related factors, such as the frequent handling of loads or handling them with high force, highly repetitive tasks, and works above the shoulder level being considered extrinsic factors leading to tendinopathy [21-23]. Raising the hand above the shoulder level is among the major causes of rotator cuff tendinopathy (RCT) among workers [22, 24-26]. In some studies, heavy manual work [27], repetitive tasks [10, 28], high frequency of work [5, 28, 29], high-force exertion [28, 29], and vibration works [29] have been reported to be related to the increased risk of structure tendinopathy. Similarly, in a recent systematic review and a meta-analysis in 2019, DM, being over age 50, and working at the overhead level were reported to be related to the increased risk of structure tendinopathy [30]. In some studies, doing work with the shoulder placed above 90° was shown to be related to the increased risk of structure tendinopathy among the working population [24, 26, 27]. However, other mechanical workplace factors, except heavy manual works [27], repetitive works [10, 28], high frequency of work [5, 28, 29], high force exertion [29, 31], and vibration work [29,31] have not been clearly associated with the increased risk of structure tendinopathy. In 1997, the National Institute for Occupational Safety and Health

published a critical synopsis of evidence-based medical investigations into the relationship between work activities and the development of MSDs. Accordingly, there was clear evidence that definite levels of static contraction, prolonged static loads, and awkward postures involving the neck and shoulder muscles were related to an increased risk of MSDs [32]. In fact, it seems that the effect of cumulative trauma is usually raised in this context, with occupational risk factors being often cumulative over time, possibly increasing the risk of shoulder diseases and injuries. Three mechanisms are introduced to provide explanations for the occurrence of cumulative trauma disorders (CTDs), which include (1) repetitive mechanical irritation, (2) micro ischemia, and (3) accumulation of metabolic products that promote inflammations or interfere with neuromuscular function [33-36]. One or more of the activities, such as excessive force, repetitive movements, awkward postures, prolonged static postures, and vibration have been associated with CTDs and RCT [37]. In most studies, only a limited number of risk factors have been considered for research on occupational risk factors in shoulder injuries. Thus, cumulative effects of these risk factors and their role have not been evaluated. Additionally, most cases of MSDs have been reported by self-reported questionnaires, which can reduce the accuracy of any association between occupational risk factors and the prevalence of MSDs. In this study, radiological findings could confirm the existence of shoulder abnormalities. Therefore, this study aims to investigate cumulative effects of risk factors on RCT among workers.

Materials and Methods

Initial data of this cross-sectional study were extracted from the full archive of the MRI imaging of Rassol-e- Akram Hospital in Tehran, Iran, for the 9-month period from July 2018 to March 2019. Accordingly, the data included MRI images of the referred patients from orthopedic and rheumatology wards to the radiology department, who had neck and shoulder pain and were suspected of rotator cuff tendinopathy (n = 470). The exclusion criteria were patients less than 18 or over 60 years old, patients whose occupation was not determined, professional athletes and patients with a history of multiple trauma, as well as patients with degenerative intervertebral disc disorders, congenital spinal disorders, malignancies, fractures or spinal surgeries in the last six months (n = 64). Finally, 406 patients remained after excluding 64 patients (n = 406).

Next, the included patients were divided into two groups based on their rotator cuff involvement, with the first group being without tendinopathy (n = 112) and the second group being involved with it (n = 294). The study protocol was reviewed and approved by the Ethics Committee of Iran University of Medical Sciences (code: IR.IUMS.FMD.REC 1396.9511308004).

The study participants were interviewed in person. The first part of the interview included general information on the participants' age, sex, education level, marital status, being a smoker or a passive smoker, body mass index, history of underlying diseases, medication use, and shoulder trauma. Besides, the second part of the interview included occupational information on their job title, work experience, employment status, daily and weekly working hours, shift work status, number of monthly shift works, and job satisfaction. Additionally, a question was asked to assess the patients' job satisfaction, and they were asked to rate their job satisfaction on a 1-10 scale.

Ergonomic risk factors were explained to the study participants, and they were asked to mark the risk factors to which they were exposed, including excessive hand raising (an arm above the shoulder > 2h/day), repetitive tasks, upper arm flexion, grip force, awkward postures, vibration, as well as shoulder abduction and rotation.

To determine cumulative effects of biomechanical risk factors altogether, the subjects were divided into several categories according to the number of the affecting biomechanical risk factors. In the first, second, third, and fourth models, people had at least 1, 2, 3, and 4 affecting biomechanical risk factors, respectively.

The study data were analyzed by SPSS V24.0. For quantitative data, mean and standard deviation were used, and for qualitative data, frequency and percentage were used. For the statistical analysis of quantitative and qualitative data, an independent sample t-test and a chi-square test were used, respectively. In addition, univariate and multivariate analyses were performed using the

Cox regression model to identify predictors of rotator cuff tendinopathy. Besides, all statistical test results less than 0.05 were considered significant.

Results

This cross-sectional study was performed on 406 patients suspected of rotator cuff tendinopathy, who were referred by orthopedic and rheumatology wards to the radiology department. The majority of the participants were male (n = 308, 75.89% of the persons). In addition, their mean age was 42.02 ± 12.63, their work experience was 8.72 ± 6.81 years, 380 (93.6%) individuals were married, 147 (36.2%) individuals had a high school and diploma, 259 (63.8%) individuals had a higher education, 106 (26.1%) individuals were smokers, 235 (58%) individuals would perform physical work, 137 (33.7%) individuals would do a shift work, 60 (14.8%) individuals were night shift workers, and 228 (56.2%) individuals felt no job satisfaction. According to radiologic findings, 294 (72%) patients had rotator cuff tendinopathy, and 112 (28%) individuals did not have tendinopathy. In patients with tendinopathy, 149 (36.69%) patients had partial width and partial tear, 149 (36.69%) patients had partial width and partial tear, 80 (19.70%) patients had complete thickness tear, 185 (45.56%) patients had tendinosis, 31 (7.63%) patients had complete tear and atrophy, 68 (16.74%) patients had the gap and complete tear, 45 (11.8%) patients had partial width, partial tear, and tendonitis, and 54 (13.3%) patients had full thickness and partial tear.

In this study, 194 (47.8%) patients had excessive hand raising (the arm above the shoulder > 2h/day), 184 (45.3) patients had repetitive tasks, 145 (35.7) patients had upper arm flexion, 112 (27.6) patients had grip force, 190 (46.8) patients had an awkward posture, 38 (9.4) patients experienced vibration, 246 (60.6) patients had shoulder abduction, and 86 (21.2) patients had shoulder rotation. Table 1 shows the participants' demographic and occupational characteristics.

Table 1. Demographic and occupational characteristics of the patients referred from orthopedic and rheumatology wards to the radiology department of Rasool-e-Akram Hospital

Variables		Mean ± SD / Number (%)
Age (year)		42.02± 12.63
Work experience (year)		8.72 ± 6.81
BMI (Kg/M ²)		25.36±2.85
Gender	Female	308 (75.89)
	Male	94 (24.14)
Marital status	Married	380 (93.6)
	Single	126 (6.4)
Smoking	Smoker	106 (26.1)
	Non-smoker	300 (73.9)

Education	High school and diploma	147 (36.2)
	Higher education	259 (63.8)
Job category	Physical	235 (58)
	Non-physical	171 (42)
Shift work	Yes	137 (33.7)
	No	269 (66.3)
Night shift	Yes	60 (14.8)
	No	346 (85.2)
Job satisfaction	Yes	178 (43.8)
	No	228 (56.2)

According to the univariate analysis, being male, being over 40 years old, and physical work significantly affected frequency rotator cuff tendinopathy ($p < 0.05$). In addition, the body mass index (BMI) of over 25 kg/m², being a smoker,

work experience of over 8 years, shift working, night shift working, and job satisfaction had no significant relationship with frequency rotator cuff tendinopathy ($p > 0.05$) (Table 2).

Table 2. Univariate and multivariate analyses of frequency rotator cuff tendinopathy in the study participants

Variables	Univariate analysis		Multivariate analysis	
	OR (95%CI)	P-value	OR (95%CI)	P-value
Age (year)	40 \geq	1	1	
	40<	2.63 (1.68 to 4.22)	0 < 001	1.12 (0.78 to 1.09)
Sex	Female	1	1	
	Male	0.77 (0.47 to 1.56)	0.03	0.38 (0.28 to 1.04)
Marital status	Married	1		
	Single	0.9 (0.81 to 1.65)	0.37	
BMI	BMI \leq 25	1		
	BMI > 25	0.6 (0.22 to 1.49)	0.80	0.94 (1.18 to 2.75)
Smoking	Non-smoker	1		
	Smoker	1.04 (0.64 to 1.71)	0.90	
Work category	Non-physical	1		
	Physical	1.25 (1.04 to 2.73)	0.02	1.1 2 (1.01 to 2.18)
Work experience	8 \geq	1		
	8<	0.77 (0.49 to 1.22)	0.16	0.58 (0.19 to 3.62)
Shift working	Non-shift worker	1		
	Shift worker	0.53 (0.28 to 2.17)	0.41	
Night shifting	Non-night shift worker	1		
	Night shift worker	0.92 (0.22 to 1.31)	0.32	
Job satisfaction	Yes	1		
	No	1.01 (0.57 to 1.92)	0.08	

According to the univariate analysis, biomechanical risk factors, such as excessive hand rising (OR=2.73 95%CI: 1.7-4.3), repetitive tasks (OR=4.37 95%CI: 2.68-7.26), upper arm flexion (OR=4.69 95%CI: 2.63-8.35), and awkward postures (OR=2.08 95%CI: 1.3-3.2) had a significant relationship with the frequency of rotator cuff tendinopathy; however, risk factors, including rotation (grip force, vibration, and abduction) had no significant relationship with the frequency of rotator cuff tendinopathy ($p > 0.05$).

According to the multivariate analysis and using the Cox regression model, risk factors, such as excessive hand raising, upper arm flexion, repetitive tasks, and awkward postures had greater effects on rotator cuff tendinopathy than sex ($p = 0.74$), age ($p = 0.09$), BMI ($p = 0.15$), and work experience ($p = 0.2$), which had no significant association. Tables 2 and 3 show the results of the univariate and multivariate analyses using the Cox regression model.

Table 3. Univariate and multivariate analyses of frequency rotator cuff tendinopathy in the study participants

Variables	Univariate analysis		Multivariate analysis		
		OR (95%CI)	P-value	OR (95%CI)	P-value
Excessive hand raising	No	1		1	
	Yes	2.73 (1.71 to 4.32)	< 0.001	2.13 (1.92 to 3.64)	0.000
Repetitive tasks	No	1		1	
	Yes	4.37 (2.68 to 7.26)	< 0.001	3.18 (1.58 to 3.79)	0.000
Upper arm flexion	No	1		1	
	Yes	4.69 (2.63 to 8.35)	< 0.001	2.19 (1.54 to 2.87)	0.000
Grip force	No	1		1	
	Yes	0.78 (0.48 to 1.25)	0.32		
Awkward posture	No	1		1	
	Yes	2.08 (1.34 to 3.25)	0.001	1.98 (1.05 to 2.17)	0.000
Vibration	No	1		1	
	Yes	0.80 (0.39 to 1.66)	0.57		
Shoulder abduction	No	1		1	
	Yes	0.54 (0.34 to 0.87)	0.21		
Shoulder rotation	No	1		1	
	Yes	1.33 (0.76 to 2.32)	0.34		

In this study, 15% (n = 61) of the participants were not affected by biomechanical risk factors, 29.8% (n = 121) were affected by one biomechanical risk factor, 27.1% (n = 110) were affected by two biomechanical risk factors, 20.4% (n = 83) were affected by three biomechanical risk factors, and 7.6% (n = 13) were affected by four biomechanical risk factors. Additionally, there was a statistically significant relationship between experiencing at least one affecting risk factor and shoulder tendinopathy (p=0<000 OR=3.96 95%CI: 2.26-6.96, experiencing at least two affecting risk factors and shoulder tendinopathy (p=0<000 OR=6.82 95%CI: 4.13-11.26), experiencing at least three risk factors and shoulder tendinopathy (p=0<000 OR=10.25 95%CI: 4.35-24.14), and experiencing at least four risk factors and shoulder tendinopathy (p=0<000 OR=12.61 95%CI: 1.69-93.63).

Discussion

In this study, conducted on 406 patients suspected of rotator cuff tendinopathy, 72% of the patients had different forms of rotator cuff abnormality. The results of this study showed that the male sex, being over 40 years old, physical work, and biomechanical risk factors, such as excessive hand raising, repetitive tasks, upper arm flexion, and awkward postures significantly affected frequency rotator cuff tendinopathy. Moreover, rotator cuff tendinopathy had no significant relationship with the participants' body mass index (BMI), smoking, work experience of over 8 years, shift working, night shift working, job satisfaction, and biomechanical risk factors, including rotation, grip force, and vibration.

In this study, RCT was higher in men than in women. Consistent with the present study, RCT was reported to be more in men than in women in some studies [4, 6]; however, there was no difference between sexes in developing rotator cuff in others studies [9, 20, 24]. This gender difference probably reflects differences in work exposure where men usually do more physical work than women. In the present study, job-related factors differed by gender. This gender difference probably reflects differences in exposure to work constraints. According to several studies, job content is different between men and women, even when the job title is the same [29, 34]. RCT has multifactorial causes, with extrinsic and intrinsic factors, or a combination of both involved [19]. Our findings showed that in workers over age 45, RCT was more common; thus, age appears to be a significant factor in the etiology of rotator cuff tendinopathy among the working population, with this being due to age-related degenerative changes [10, 24, 35, and 36].

As a person ages, they will be more likely to be exposed to occupational risk factors, with the prevalence of shoulder involvement increased, as in some studies, RCT increased with age [4, 28]. Decreased ultimate strain, decreased ultimate load, decreased elasticity, and decreased overall tensile strength have been reported in some studies in aging tendons [38, 39]. In addition, microscopic and biochemical pathological changes have led to the degeneration of tenocytes and collagen fibers as well as accumulation of lipids and ground substance (glycosaminoglycans) in aging tendons [35].

However, there is no consensus on whether pathological changes in rotator cuff tendons are primarily due to aging or they are the secondary consequence of excessive mechanical factors and overload [19].

In the present study, no relationship was found between the BMI and rotator cuff involvement. Results from a study [24] were consistent with ours, but they were inconsistent with the results obtained from another study [4].

In this study, excessive hand raising, repetitive tasks, upper arm flexion, and awkward postures were the most important mechanical risk factors for RCT among the workers. These results of our study were consistent with those of a number of other studies. In some studies [22, 24, and 40], hand-up work with the shoulder being over 90° was revealed to be associated with an increased risk of rotator cuff tendinopathy among the working population. In our study, arm abduction had no relationship with rotator cuff tendinopathy, but it had a relationship in a study [24]. In our study, vibration had no association with RCT, but in some studies, the opposite was true [5, 29]. The results of the four models used in the present study showed that the incidence of tendonitis increased with an increase in the number of mechanical risk factors in the workplace. Based on model 1, having at least one affecting risk factor, two affecting risk factors, three affecting risk factors, and four affecting risk factors increased the chances of developing RCT in workers by 3.96, 6.82, 10.25, and 12.61 times, respectively.

In fact, occupational risk factors often become cumulative over time, thereby increasing the danger of shoulder diseases and injuries. In most workplaces, there is usually over one mechanical risk factor, so it is necessary to pay attention to the cumulative effects of all of them.

Several limitations existed in the present study. Firstly, it was a cross-sectional study, so we could not determine the cause-and-effect relationships; secondly, we categorized patients' jobs into physical and non-physical jobs, and perhaps it would have been better to go into more detail; thirdly, we were unable to investigate psychological factors effecting CRT. Therefore, it is suggested that a larger sample size be selected in the form of a cohort study to take into account more details.

This study was one of the first ones conducted in Iran at a major radiological center to investigate cumulative effects of risk factors on RCT among workers, in which, we evaluated ergonomic risk factors along with other individual risk factors.

Conclusion

The results of this study showed that age over 45, excessive hand raising, repetitive tasks, upper arm flexion, and awkward postures were significantly associated with RCT, with the accompanying occupational risk factors increasing the risk of RCT in the workplace. Based on the findings of this study, the existence of over one mechanical risk factor in the workplace can be associated with a higher increase in RCT. In most workplaces, there is usually over one mechanical risk factor, so it is necessary to pay attention to the cumulative effects of all of them.

Acknowledgement

We would like to extend our thanks to the deputy for research at Iran University of Medical Sciences for his assistance in doing this research (code IR.IUMS.FMD.REC 1396.9511308004).

Conflict of interest: None declared.

References

1. Luime JJ, Koes BW, Hendriksen IJ, Burdorf A, Verhagen AP, Miedema HS, et al. Prevalence and incidence of shoulder pain in the general population; a systematic review. *Scand J Rheumatol.* 2004;33(2):73-81.
2. Hill CL, Gill TK, Shanahan EM, Taylor AW. Prevalence and correlates of shoulder pain and stiffness in a population-based study: the North West Adelaide Health Study. *Int J Rheum Dis.* 2010;13(3):215-22.
3. Linsell L, Dawson J, Zondervan K, Rose P, Randall T, Fitzpatrick R, et al. Prevalence and incidence of adults consulting for shoulder conditions in UK primary care; patterns of diagnosis and referral. *Rheumatology.* 2005;45(2):215-21.
4. Miranda H, Punnett L, Viikari-Juntura E, Heliövaara M, Knekt P. Physical work and chronic shoulder disorder. Results of a prospective population-based study. *Ann Rheum Dis.* 2008;67(2):218-23.
5. Silverstein BA, Bao SS, Fan ZJ, Howard N, Smith C, Spielholz P, et al. Rotator cuff syndrome: personal, work-related psychosocial and physical load factors. *J Occup Environ Med.* 2008;50(9):1062-76.
6. Silverstein BA, Viikari-Juntura E, Fan ZJ, Bonauto DK, Bao S, Smith C. Natural course of nontraumatic rotator cuff tendinitis and shoulder symptoms in a working population. *Scand J Work Environ Health.* 2006;32(2):99-108.
7. Kaergaard A, Andersen JH. Musculoskeletal disorders of the neck and shoulders in female sewing machine operators: prevalence,

- incidence, and prognosis. *Occup Environ Med.* 2000;57(8):528-34.
8. Bonde JP, Mikkelsen S, Andersen JH, Fallentin N, Baelum J, Svendsen SW, et al. PRIM Health Study Group. Prognosis of shoulder tendonitis in repetitive work: a follow up study in a cohort of Danish industrial and service workers. *Occup Environ Med.* 2003;60(9):E8.
 9. Walker-Bone K, Palmer KT, Reading I, Coggon D, Cooper C. Prevalence and impact of musculoskeletal disorders of the upper limb in the general population. *Arthritis Rheum.* 2004;51(4):642-51.
 10. Roquelaure Y, Bodin J, Ha C, Petit Le Manac'h A, Descatha A, Chastang JF, et al. Personal, biomechanical, and psychosocial risk factors for rotator cuff syndrome in a working population. *Scand J Work Environ Health.* 2011;37(6):502-11.
 11. Lewis J, McCreesh K, Roy JS, Ginn K. Rotator Cuff Tendinopathy: Navigating the Diagnosis-Management Conundrum. *J Orthop Sports Phys Ther.* 2015;45(11):923-37.
 12. Desmeules F, Boudreault J, Dionne CE, Fremont P, Lowry V, MacDermid JC, et al. Efficacy of exercise therapy in workers with rotator cuff tendinopathy: a systematic review. *J Occup Health.* 2016;58(5):389-403.
 13. Biau DJ, Porcher R. Letter to the editor re: Orthopaedic surgeons prefer to participate in expertise-based randomized trials: Bednarska E, Bryant D, Devereaux, PJ. Orthopaedic surgeons prefer to participate in expertise-based randomized trials. *Clin Orthop Relat Res.* 2008;466:1734-1744. *Clin Orthop Relat Res.* 2009;467(1):298-300; author reply 1-2.
 14. Croft P, Pope D, Silman A. The clinical course of shoulder pain: prospective cohort study in primary care. Primary Care Rheumatology Society Shoulder Study Group. *BMJ.* 1996;313(7057):601-2
 15. van der Windt DA, Koes BW, de Jong BA, Bouter LM. Shoulder disorders in general practice: incidence, patient characteristics, and management. *Ann Rheum Dis.* 1995;54(12):959-64.
 16. Brox JI, Gjengedal E, Uppheim G, Bohmer AS, Brevik JI, Ljunggren AE, et al. Arthroscopic surgery versus supervised exercises in patients with rotator cuff disease (stage II impingement syndrome): a prospective, randomized, controlled study in 125 patients with a 2 1/2-year follow-up. *J Shoulder Elbow Surg.* 1999;8(2):102-11.
 17. Holmgren T, Hallgren HB, Oberg B, Adolfsson L, Johansson K. Effect of specific exercise strategy on need for surgery in patients with subacromial impingement syndrome: randomised controlled study. *Br J Sports Med.* 2014;48(19):1456-7.
 18. Ludewig PM, Borstad JD. Effects of a home exercise programme on shoulder pain and functional status in construction workers. *Occup Environ Med.* 2003;60(11):841-9.
 19. Seitz AL, McClure PW, Finucane S, Boardman ND 3rd, Michener LA. Mechanisms of rotator cuff tendinopathy: intrinsic, extrinsic, or both? *Clin Biomech (Bristol, Avon).* 2011;26(1):1-12.
 20. Soslowky LJ, Thomopoulos S, Esmail A, Flanagan CL, Iannotti JP, Williamson JD, 3rd, et al. Rotator cuff tendinosis in an animal model: role of extrinsic and overuse factors. *Ann Biomed Eng.* 2002;30(8):1057-63.
 21. Rechartdt M, Shiri R, Karppinen J, Jula A, Heliövaara M, Viikari-Juntura E. Lifestyle and metabolic factors in relation to shoulder pain and rotator cuff tendinitis: a population-based study. *BMC Musculoskelet Disord.* 2010;11:165.
 22. Miranda H, Viikari-Juntura E, Heistaro S, Heliövaara M, Riihimäki H. A population study on differences in the determinants of a specific shoulder disorder versus nonspecific shoulder pain without clinical findings. *Am J Epidemiol.* 2005 ;161(9):847-55.
 23. Viikari-Juntura E, Shiri R, Solovieva S, Karppinen J, Leino-Arjas P, Varonen H, et al. Risk factors of atherosclerosis and shoulder pain--is there an association? A systematic review. *Eur J Pain.* 2008;12(4):412-26.
 24. Bodin J, Ha C, Petit Le Manac'h A, Sérazin C, Descatha A, Leclerc A, et al. Risk factors for incidence of rotator cuff syndrome in a large working population. *Scand J Work Environ Health.* 2012;38(5):436-46.
 25. Grzywacz JG, Arcury TA, Mora D, Anderson AM, Chen H, Rosenbaum DA, et al. Work organization and musculoskeletal health: clinical findings from immigrant Latino poultry processing and other manual workers. *J Occup Environ Med.* 2012;54(8):995-1001.
 26. Svendsen SW, Bonde JP, Mathiassen SE, Stengaard-Pedersen K, Frich LH. Work related shoulder disorders: quantitative exposure-response relations with reference to arm posture. *Occup Environ Med.* 2004 Oct;61(10):844-53
 27. Mehta SK, Teefey SA, Middleton W, Steger-May K, Sefko JA, Keener JD. Prevalence and risk factors for development of subscapularis and biceps pathology in shoulders with degenerative rotator cuff disease: a prospective cohort evaluation. *J Shoulder Elbow Surg.* 2020;29(3):451-8
 28. Frost P, Bonde JP, Mikkelsen S, Andersen JH, Fallentin N, Kaergaard A, et al. Risk of shoulder tendinitis in relation to shoulder loads in monotonous repetitive work. *Am J Ind Med.* 2002;41(1):11-8.
 29. Silverstein B, Fan ZJ, Smith CK, Bao S, Howard N, Spielholz P, et al. Gender adjustment or stratification in discerning upper extremity musculoskeletal disorder risk? *Scand J Work Environ Health.* 2009;35(2):113-26.
 30. Leong HT, Fu SC, He X, Oh JH, Yamamoto N,

- Hang S. Risk factors for rotator cuff tendinopathy: A systematic review and meta-analysis. *J Rehabil Med.* 2019;51(9):627-37.
31. Sutinen P, Toppila E, Starck J, Brammer A, Zou J, Pyykkö I. Hand-arm vibration syndrome with use of anti-vibration chain saws: 19-year follow-up study of forestry workers. *Int Arch Occup Environ Health.* 2006;79(8):665-7.
32. Solomonow M. Sensory-motor control of ligaments and associated neuromuscular disorders. *J Electromyogr Kinesiol.* 2006;16(6):549-67.
33. Barbe MF, Barr AE. Inflammation and the pathophysiology of work-related musculoskeletal disorders. *Brain Behav Immun.* 2006 Sep;20(5):423-9.
34. de Zwart BC, Frings-Dresen MH, Kilbom A. Gender differences in upper extremity musculoskeletal complaints in the working population. *Int Arch Occup Environ Health.* 2001;74(1):21-30.
35. Kannus P, Jozsa L. Histopathological changes preceding spontaneous rupture of a tendon. A controlled study of 891 patients. *J Bone Joint Surg Am.* 1991;73(10):1507-25.
36. Teunis T, Lubberts B, Reilly BT, Ring D. A systematic review and pooled analysis of the prevalence of rotator cuff disease with increasing age. *J Shoulder Elbow Surg.* 2014;23(12):1913-21.
37. Miranda H, Punnett L, Viikari-Juntura E, Heliövaara M, Knekt P. Physical work and chronic shoulder disorder. Results of a prospective population-based study. *Ann Rheum Dis.* 2008;67(2):218-23.
38. Best TM, Kirkendall DT, Almekinders LC, Garrett WE. Basic Science and Injury of Muscle, Tendon, and Ligaments. In: DeLee JD, Drez D, Miller MD, editors. *Orthopaedic sports medicine: principles and practice.* Philadelphia, United States: W.B. Saunders; c2003.
39. Woo SL. Anatomy, biology, and biomechanics of tendon and ligament. In: Buckwalter JA, Einhorn TA, Simon SR, American Academy of Orthopaedic Surgeons, editors. *Orthopaedic basic science: biology and biomechanics of the musculoskeletal system.* 2nd ed. Rosemont, IL, United States: American Academy of Orthopaedic Surgeons; 2000. P.581-616.
40. Andersen JH, Haahr JP, Frost P. Risk factors for more severe regional musculoskeletal symptoms: a two-year prospective study of a general working population. *Arthritis Rheum.* 2007;56(4):1355-64.